Alcom-FT Technical Report Series
ALCOMFT-TR-02-19

Rapidly Mixing Markov Chains for Sampling Contingency Tables with a Constant Number of Rows *

Mary Cryan
School of Computing
University of Leeds
Leeds LS2 9JT, United Kingdom
maryc@comp.leeds.ac.uk

Martin Dyer
School of Computing
University of Leeds
Leeds LS2 9JT, United Kingdom
dyer@comp.leeds.ac.uk

Leslie Ann Goldberg
Department of Computer Science
University of Warwick
Coventry CV4 7AL, United Kingdom
leslie@dcs.warwick.ac.uk

Mark Jerrum
Division of Informatics
University of Edinburgh
Edinburgh EH9 3JZ, United Kingdom
mrj@dcs.ed.ac.uk

Russell Martin
Department of Computer Science
University of Warwick
Coventry CV4 7AL, United Kingdom
martin@dcs.warwick.ac.uk

May 2, 2002

Abstract

We consider the problem of sampling almost uniformly from the set of contingency tables with given row and column sums, when the number of rows is a constant. Cryan and Dyer [2] have recently given a fully polynomial randomized approximation scheme (*fpras*) for the related counting problem, which only employs Markov chain methods indirectly. But they leave open the question as to whether a natural Markov chain on such tables mixes rapidly. Here we answer this question in the affirmative, and hence provide a very different proof of the main result of [2]. We show that the "2×2 heat-bath" Markov chain is rapidly mixing. We prove this by considering first a heat-bath chain operating on a larger window. Using techniques developed by Morris [17] for the multidimensional knapsack problem, we show that this chain mixes rapidly. We then apply the comparison method of Diaconis and Saloff-Coste [6] (see also Randall and Tetali [21]) to show that the 2×2 chain is rapidly mixing. To establish this, we provide the first proof that this chain mixes in time polynomial in the input size even when both the number of rows and number of columns are constant.

*This work was partially supported by the EPSRC grant "Sharper Analysis of Randomised Algorithms: a Computational Approach", the EPSRC grant GR/R44560/01 "Analysing Markov-chain based random sampling algorithms" and the IST Programme of the EU under contract numbers IST-1999-14186 (ALCOM-FT) and IST-1999-14036 (RAND-APX).

1 Introduction

Given two vectors of positive integers, $r = (r_1, \ldots, r_m)$ and $c = (c_1, \ldots, c_n)$, an $m \times n$ matrix $[X_{i,j}]$ of non-negative integers is a *contingency table* with row sums r and column sums c if $\sum_{j=1}^{n} X_[i,j] = r_i$ for every row i and $\sum_{i=1}^{m} X_[i,j] = c_j$ for every column j. We write $\Sigma_{r,c}$ to denote the set of all contingency tables with row sums r and column sums c. We assume that $\sum_{i=1}^{m} r_i = \sum_{j=1}^{n} c_j$ (since otherwise $\Sigma_{r,c} = \emptyset$) and denote by N the common total, called the *table sum*.

In this paper, we consider the problem of sampling contingency tables almost uniformly at random. Unfortunately, no technique currently exists for polynomial time sampling of general contingency tables with arbitrary row and column sums. Here we consider a particular restriction, the case where the number of rows is a constant. Cryan and Dyer [2] recently gave a fully polynomial randomized approximation scheme for *counting* contingency tables in this setting (i.e. for approximating $|\Sigma_{r,c}|$ with given r, c). Although the problem is not self-reducible, a variant of the method of [15] can be used to show equivalence of approximate counting and sampling. (It is known that the problem of *exact* counting is #P-Complete [10].) The algorithm of [2] is a mixture of dynamic programming and volume estimation, and uses Markov chain methods only indirectly. So [2] leaves open the question as to whether the Markov chain Monte Carlo (MCMC) method can be applied *directly* to this problem. In addition to its intrinsic interest, this question has importance for two reasons. Firstly, previous research in this area has routinely adopted the MCMC approach. Secondly, the MCMC method is more convenient, and has been more widely applied, for practical applications of sampling.

We give here the first proof of rapid mixing for a natural Markov chain when the number of rows m is a constant. Our final chain, the chain of Dyer and Greenhill [9], randomly modifies a uniformly selected 2×2 subtable. However, we first analyse a chain which randomly modifies a $m \times (2d_m + 1)$ subtable, where d_m is a constant we will describe later. We use the multicommodity flow technique of Sinclair [22] to analyse the mixing time of this $m \times (2d_m + 1)$ "heat-bath" chain. Using techniques developed by Morris [17] (building on work of Morris and Sinclair [18]), we are able to show that this chain mixes in time polynomial in the number of rows, the number of columns, and the logarithm of the table sum. In Section 4 we compare this chain to the Dyer-Greenhill chain (2×2 heat-bath) and hence show that the Dyer-Greenhill chain is also rapidly mixing. It may be observed that no proof was previously known that the Dyer-Greenhill (or any other) chain converges in polynomial time even when the number of columns, as well as the number of rows, is constant. Establishing this fact is one step of our proof. (See Pak [20] for an approach to this problem not using MCMC.) We note further that our results provide an alternative (and very different) *fpras* for this problem to that of Cryan and Dyer[2].

Contingency tables are important in applied statistics, where they are used to summarize the results of tests and surveys. The conditional volume test of Diaconis and Efron [4] is perhaps the most soundly based method for performing tests of significance in such tables. The Diaconis-Efron test provides strong motivation for the problem of efficiently choosing a contingency table with given row and column sums uniformly at random. Other applications of counting and sampling contingency tables are discussed by Diaconis and Gangolli [5]. See also Mount [19] for additional information, and De Loera and Sturmfels [3] for the current limits of exact counting methods.

With the notable exception of [2], most previous work on sampling contingency tables applies the MCMC method, as described in the survey of Jerrum and Sinclair [14]. This method, which has been used to solve many different sampling problems, is based on a very simple idea. Suppose that we have a Markov chain on a finite set of discrete structures Ω, defined by the transition matrix P. If the Markov chain is *ergodic*, then it will converge to a unique stationary distribution π on Ω, regardless of the initial state. This gives a nice method for sampling from the distribution π: starting in any state, we run the Markov chain for some "sufficiently long" number of steps. Then the final state is taken as a sample from π. The key issue with using the MCMC method is determining how long the chain takes to converge to its stationary distribution.

The first explicit definition of Markov chains for uniformly sampling contingency tables apparently occurs in the papers of Diaconis and Gangolli [5] and Diaconis and Saloff-Coste [7], although it is mentioned in [5] that this chain had already been used by practitioners. A single step of the chain is generated as follows: an ordered pair of rows i_1, i_2 are chosen uniformly at random from all rows of the table, and an ordered pair of columns j_1, j_2 are chosen uniformly at random from all columns, giving a 2×2 submatrix. The entries of

the 2×2 submatrix are modified as follows:

$$X'[i_1, j_1] = X[i_1, j_1] + 1 \qquad X'[i_1, j_2] = X[i_1, j_2] - 1$$
$$X'[i_2, j_1] = X[i_2, j_1] - 1 \qquad X'[i_2, j_2] = X[i_2, j_2] + 1$$

If modifying the matrix results in a negative value for any $X'[i, j]$, the move is not carried out. Diaconis and Gangolli proved that this Markov chain is ergodic, and the stationary distribution of the chain is uniform on $\Sigma_{r,c}$. They did not attempt to bound the mixing time of the chain, but it is clear that the mixing time can never be better than pseudopolynomial in the input.

Later Diaconis and Saloff-Coste [7] considered the case when the numbers of rows and columns are both constant and proved that, in this case, their chain converges in time quadratic in the table sum. Hernek [13] considered the case when the table has two rows and proved that the same chain mixes in time polynomial in the number of columns and the table sum. Chung et al. [1] showed that a slightly modified version of the Diaconis and Saloff-Coste chain converges in time polynomial in the table sum, the number of rows and the number of columns, provided that all row and column sums are sufficiently large.

The first truly polynomial-time algorithm (polynomial in the number of rows, the number of columns and the logarithm of the table sum) for sampling contingency tables was given by Dyer, Kannan and Mount [10]. They took a different approach to the sampling problem, considering $\Sigma_{r,c}$ as the set of integer points within a convex polytope. They used an existing algorithm for sampling continuously from a convex polytope, combined with a rounding procedure, to sample integer points from inside the polytope. For any input with row sums of size $\Omega(n^2 m)$ and column sums of size $\Omega(nm^2)$, their algorithm converges to the uniform distribution on $\Sigma_{r,c}$ in time polynomial in the number of rows, the number of columns, and the logarithm of the table sum. Their result was later refined by Morris [16], who showed that the result also holds when the row sums are $\Omega(n^{3/2} m \log m)$ and the column sums are $\Omega(m^{3/2} n \log n)$.

Using different techniques, Dyer and Greenhill [9] considered the problem of sampling contingency tables when the table has only two rows. They considered a natural 2×2 heat-bath chain. In the two-row case, a single step of the Dyer-Greenhill chain at contingency table X is performed as follows: two columns j_1, j_2 are chosen uniformly at random from all columns, giving a 2×2 submatrix (since the table only has two rows) with column sums c_{j_1}, c_{j_2} and a pair of induced row sums s_1, s_2. A 2×2 submatrix is then chosen uniformly at random from the set of all tables with the induced row sums (s_1, s_2) and column sums (c_{j_1}, c_{j_2}). The j_1th and j_2th columns of X are replaced by this new subtable. Dyer and Greenhill showed that for two-rowed tables, their chain converges to the uniform distribution on $\Sigma_{r,c}$ in time that is polynomial in the number of columns and the logarithm of the table sum.

This paper can properly be viewed as extending Dyer and Greenhill's results to any constant number of rows. Thus, our main result is that the 2×2 heat-bath chain is rapidly mixing for any constant number of rows. First, however, in Section 3, we examine the following heat-bath chain, which is a natural generalization of the Dyer-Greenhill chain to tables with any constant number of rows m. For each m we will define a constant d_m which depends only on m. A step of the chain, starting from a contingency table, X, is as follows. First, $(2d_m + 1)$ columns $j_1, \ldots, j_{2d_m+1}$ are chosen uniformly at random from the columns of X. Then the induced row sums $s_1, \ldots, s_m$ are calculated and the chosen columns of X are replaced with a subtable selected uniformly from all tables with row sums $s_1, \ldots, s_m$ and column sums $c_{j_1}, \ldots, c_{j_{2d_m+1}}$. We will refer to this chain as $\mathcal{M}_{\mathrm{HB}}$. The fact that a step of the chain can be carried out in polynomial time follows from [20]. Theorem 8 shows that this chain is rapidly mixing. This, together with Theorem 9, gives the main result.

2 Background

In this section we summarize the techniques that we will use to bound the mixing time of our heat-bath chain. Our analysis is carried out using the multicommodity flow approach of Sinclair [22] for bounding the mixing time of a Markov chain. Sinclair's result builds on some earlier work due to Diaconis and Stroock [8].

The setting is familiar: we have a finite set Ω of discrete structures, and a transition matrix P on the state space Ω. It is assumed that the Markov chain defined by P is *ergodic*, that is, it satisfies the properties of *irreducibility* and *aperiodicity* (see Grimmett and Stirzaker [11]). It is well-known that any ergodic Markov chain has a unique *stationary distribution*, that is, there is a unique distribution π on Ω such

that $\pi P = \pi$. Furthermore, for any choice of initial state $x \in \Omega$ and any state $y \in \Omega$, $P^t(x,y) \to \pi(y)$ as $t \to \infty$ (see page 208 of Grimmett and Stirzaker [11] for details). Sinclair also assumes that the Markov chain is *reversible* with respect to its stationary distribution, that is, $\pi(x)P(x,y) = \pi(y)P(y,x)$ for all $x, y \in \Omega$.

For any start state x, we define the *variation distance* between the stationary distribution and a walk of length t by $\mathrm{V}(\pi, P^t(x)) = (1/2)\sum_{y\in\Omega} |\pi(y) - P^t(x,y)|$. For any $0 < \epsilon < 1$ and any start state x, let $\tau_x(\epsilon)$ be defined as $\tau_x(\epsilon) = \min\{t : \mathrm{V}(\pi, P^t(x)) \le \epsilon\}$. The *mixing time* of the chain is given by the function $\tau(\epsilon)$, defined as $\tau(\epsilon) = \max\{\tau_x(\epsilon) : x \in \Omega\}$.

The following theorem of Diaconis and Stroock [8] gives an upper bound on the mixing time of a Markov chain in terms of the second eigenvalue of the transition matrix and the stationary distribution π:

Theorem 1 (Diaconis and Stroock [8]) *Let P be the transition matrix of an ergodic reversible Markov chain with stationary distribution π. Assume that $P(x,x) \ge 1/2$ for all $x \in \Omega$, so the eigenvalues of P are all non-negative. Let $1 = \lambda_0 > \lambda_1 \ge \lambda_2 \ge \ldots \lambda_{|\Omega|-1} \ge 0$ be these eigenvalues. Then for any initial state $x \in \Omega$,*

$$\begin{aligned} \tau_x(\epsilon) &\le (\ln \pi(x)^{-1} + \ln \epsilon^{-1})/(\ln \lambda_1^{-1}) \\ &\le (\ln \pi(x)^{-1} + \ln \epsilon^{-1})/(1 - \lambda_1) \end{aligned}$$

Therefore, to prove an upper bound on the mixing time of a Markov chain, it is sufficient to find an upper bound on the eigenvalue λ_1. The multicommodity flow approach is defined in terms of a graph G_Ω defined by the Markov chain. The vertices of G_Ω are the elements of Ω, and the graph contains an edge $(u \to v)$ for every pair of states such that $P(u,v) > 0$. For any $x, y \in \Omega$, a *unit flow* from x to y is a set $\mathcal{P}_{x,y}$ of simple directed paths of G_Ω from x to y, such that each path $p \in \mathcal{P}_{x,y}$ has a positive weight α_p, and the sum of the α_p over $p \in \mathcal{P}_{x,y}$ is 1. A *multicommodity flow* is a family of unit flows $\mathcal{F} = \{\mathcal{P}_{x,y} : x, y \in \Omega\}$ containing a unit flow for every pair of states from Ω. The important properties of a multicommodity flow are the maximum flow passing through any edge and the maximum length of a path in the flow. We define the length $\mathcal{L}(\mathcal{F})$ of the multi-commodity flow $\mathcal{F}$ by $\mathcal{L}(\mathcal{F}) = \max_{x,y} \max\{|p| : p \in \mathcal{P}_{x,y}\}$, where $|p|$ denotes the length of p. For any edge e of G_Ω, we define $\mathcal{F}(e)$ to be the sum of the α_p weights over all p such that $e \in p$ and $p \in \mathcal{P}_{x,y}$ for some $x, y \in \Omega$.

The following theorem is an amalgamation of the results of Sinclair [22]:

Theorem 2 (Sinclair [22]) *Let P be the transition matrix of an ergodic, reversible Markov chain on Ω whose stationary distribution is the uniform distribution. Let $\mathcal{F}$ be a multicommodity flow on the graph G_Ω. Then the mixing time of the chain is bounded above by*

$$\tau(\epsilon) \le 2|\Omega|^{-1}\mathcal{L}(\mathcal{F}) \max_e \frac{\mathcal{F}(e)}{P(e)} (\log|\Omega| + \log \epsilon^{-1}) \tag{1}$$

Two key ingredients of our analysis of the large heat-bath chain in Section 3 are the "balanced almost-uniform permutations" and the "strongly balanced permutations" used by Morris [17] for his analysis of the multidimensional knapsack problem. We will use an interleaving of a balanced almost-uniform permutation and a strongly balanced permutation to spread flow between each pair of states $x, y \in \Sigma_{r,c}$. The main idea is this: Given x and y we will use a permutation π of the columns of x to define a path of contingency tables from x to y. We will route flow from x to y along this path. Actually, π will be chosen from a distribution and the amount of flow routed along the path corresponding to π will be proportional to the probability with which π is generated.

Definition 3 (Morris [17] Definition 2.3) *Let σ be a random variable taking values in S_n (i.e., σ is a permutation of $\{1,\ldots,n\}$) and let $\lambda \in \mathbf{R}$. Then σ is a λ-uniform permutation if*

$$\Pr[\sigma\{1,\ldots,k\} = U] \le \lambda \times \binom{n}{k}^{-1}$$

for every k with $k \in [n]$ and every $U \subseteq \{1,\ldots,n\}$ of cardinality k.

We will use the following notation. If σ is a permutation of the n columns of a contingency table, $\sigma(i)$ will denote the original column (in $[1,\ldots,n]$) which gets put in position i by the permutation. Thus, $\sigma\{1,\ldots,k\}$ above denotes the set of original columns that get put in the first k positions by the permutation.

Definition 4 (Morris [17] Definition 3.1) *Let $w_1, \ldots, w_n \in \mathbf{R}^d$ be any d-dimensional* weights *satisfying $\sum_{j=1}^n w_j = 0$ (i.e. $\sum_{j=1}^n w_j^i = 0$ for every $i \in [d]$). A permutation σ of $1, \ldots, n$ is ℓ-balanced if*

$$|\sum_{j=1}^{k} w_{\sigma(j)}^i| \quad \leq \quad \ell M_i$$

for all $i \in [d]$ and $k \in [n]$, where $M_i = \max_{1 \leq j \leq n} |w_j^i|$.

Definition 5 (Morris [17] Definition 3.3) *Let $w_1, \ldots, w_n \in \mathbf{R}^d$ be any d-dimensional weights. Define $\mu = (\mu^1, \ldots, \mu^d)$ to be the vector of means of the w_j weights ($\mu^i = (\sum_{j=1}^n w_j^i)/n$ for all i). A permutation σ of $1, \ldots, n$ is strongly ℓ-balanced if for all $k \in [n]$ and all $i \in [d]$, there exists a set $S \subseteq [n]$ with $|S \oplus \{1, \ldots, k\}| < \ell$ such that $(\sum_{j=1}^k w_{\sigma(j)}^i - k\mu^i)$ and $(\sum_{j \in S} w_{\sigma(j)}^i - k\mu^i)$ have opposite signs (or either is 0).*

Morris showed how to construct balanced almost-uniform permutations when d is constant.

Theorem 6 (Morris [17] Theorem 3.2) *For every positive integer d, there exists a constant c_d and a polynomial function p_d such that for any set of weights $\{w_j\}_{j=1}^n$ in $\mathbf{R}^d$, there exists a c_d-balanced, $p_d(n)$-uniform permutation.*

They key points which we should keep in mind are (1) the distribution which he constructs is "nearly" uniform (and has a fair amount of entropy) and (2) the permutations satisfy some sort of balance property on multi-dimensional weights. Roughly, one should think of these weights as corresponding to the columns of our contingency tables – the multiple dimensions come from having multiple rows. Morris also adapted a result of Steinitz[23] (see also Grinberg and Sevast'yanov [12]) to show that

Theorem 7 (Morris [17] Lemma 3.4) *For any sequence $\{w_j\}_{j=1}^n$ in $\mathbf{R}^d$, there exists a strongly $16d^2$-balanced permutation.*

3 Analysis of the generalized chain

Let $r = (r_1, \ldots, r_m)$ be a list of row sums and $c = (c_1, \ldots, c_n)$ a list of column sums. Let Ω be the states space $\Sigma_{r,c}$. Recall that N is the table sum $\sum_{i=1}^m r_i$. Let $d_m = 2m(3c_m + 1) + 1 + 34m^2$. Let $\mathcal{M}_{\mathrm{HB}}$ be the heat-bath Markov-chain with window-size $m \times (2d_m + 1)$ which was defined at the end of Section 1. Let P_{HB} be the transition matrix of this chain. In this section, we prove the following theorem.

Theorem 8 *The mixing time of $\mathcal{M}_{\mathrm{HB}}$ is bounded from above by a polynomial in n and N.*

In order to prove Theorem 8, we will show how to define a multicommodity flow $\mathcal{F}$ such that the total flow along any transition (ω, ω'') is at most $2fn^{2d_m+1}P_{\mathrm{HB}}(\omega, \omega'')$, where f is an expression that is at most $\mathrm{poly}(n)|\Omega|$. We will ensure that $\mathcal{L}(\mathcal{F})$ is bounded from above by a polynomial in n. Theorem 8 will then follow from (1). First, we will define a multicommodity flow $\mathcal{F}^*$ in which the total flow through any state ω is at most f. We will construct $\mathcal{F}$ by modifying $\mathcal{F}^*$. The construction of $\mathcal{F}^*$ uses the method of Morris [17].

Let k be the index of the largest column sum c_k. Let X and Y be contingency tables in Ω. Let X_j denote the jth column of X. We will show how to route a unit of flow from X to Y.

The rough idea is as follows. We will choose a permutation π from an appropriate distribution. π will be a permutation of some of the columns of the table. Actually, π will be a permutation of most of the columns of the table. The permutation π will define for us a path $X = Z'_0, \ldots, Z'_{n'+1} = Y$ where each Z'_h is in Ω and there is a transition in $\mathcal{M}_{\mathrm{HB}}$ from each Z'_h to Z'_{h+1}. The amount of flow that we route along this path will be proportional to the probability with which π is chosen.

c_m is the constant of Theorem 6 for balanced almost-uniform permutations for dimension m. Let R_i^X be the set of indices for the $3c_m + 1$ largest entries of row i of X. Let R_i^Y be the set of indices for the $3c_m + 1$ largest entries of row i of Y. Let R be the union of all the R_i^X and R_i^Y sets together with the index k. The cardinality of R is at most $2m(3c_m + 1) + 1$. The columns in R are "reserved" columns that we identify before permuting the columns. We will not permute these columns — we need them for something else. For every row i, $M_i = \min\{\max\{X[i,j] : j \notin R\}, \max\{Y[i,j] : j \notin R\}\}$. Define

$L_i = \{j : j \notin R, X[i,j] > M_i\} \cup \{j : j \notin R, Y[i,j] > M_i\}$. Note that by definition of M_i, we either have $\{j : j \notin R, X[i,j] > M_i\} = \emptyset$ or $\{j : j \notin R, Y[i,j] > M_i\} = \emptyset$. By definition, $M_i \leq X[i,j]$, for all $j \in R_i^X$, and $M_i \leq Y[i,j]$, for all $j \in R_i^Y$. $L = \cup_{i=1}^m L_i$. $S = [n] - (L \cup R)$. For every column $j \in [n] - R$, define the m-dimensional weight $w_j = Y_j - X_j$. Let μ be the m-dimensional vector representing the mean of the $w_{j \in [n]-R}$. Note that $\mu^i = (\sum_{j \in R} X[i,j] - \sum_{j \in R} Y[i,j])/(n - |R|)$.

Let π_1 be a strongly $16m^2$-balanced permutation on the set of weights $\{w_j\}_{j \in L}$. Let π_2 be a c_m-balanced $p_m(|S|)$-uniform permutation on $\{w_j\}_{j \in S}$. π_2 is a random variable. Interlacing π_1 and π_2 in the same way as Morris [17], we get a permutation π on $\{w_j\}_{j \in [n]-R}$ satisfying inequalities (3.8) and (3.9) of Morris on page 35. That is, for every prefix h (h is the index of a column), every dimension i (i is the index of a row), we have sets of column indices $V_{i,h}$ and $W_{i,h}$ such that $V_{i,h}$ differs from $\{1, \ldots, h\}$ by at most $17m^2$ indices and $W_{i,h}$ differs from $\{1, \ldots, h\}$ by at most $17m^2$ indices and

$$\sum_{j \in V_{i,h}} w^i_{\pi(j)} \leq (h-1)\mu^i + 3c_m M_i \quad \text{for every } i = 1, \ldots, m \quad (3.8)$$

$$\sum_{j \in W_{i,h}} w^i_{\pi(j)} \geq (h-1)\mu^i - 3c_m M_i \quad \text{for every } i = 1, \ldots, m \quad (3.9)$$

Let $n' = n - |R|$. We define the path of tables $X = Z_0, Z_1, ..., Z_{n'}$ as follows. For every h, Z_h contains the columns X_j for $j \in R \cup \pi\{h+1, \ldots, n'\}$ and columns Y_j for $j \in \pi\{1, \ldots, h\}$. $Z_{n'}$ differs from Y by at most $2m(3c_m + 1) + 1$ columns.

$Z_0, \ldots, Z_{n'}$ may not be contingency tables in $\sigma_{r,c}$ since they may not satisfy the row constraints. Thus, we cannot use this path as the path from X to Y. Nevertheless, we can base our path on these tables. In particular, we introduce the notation $(J(X), J(Y))$ to denote a set containing columns from X and from Y: for any $J(X) \subseteq [n]$ and $J(Y) \subseteq [n]$, $(J(X), J(Y))$ contains X_j for every $j \in J(X)$ and Y_j for $j \in J(Y)$. Let $J_h(X) = R \cup \pi\{h+1, \ldots, n'\}$ and $J_h(Y) = \pi\{1, \ldots, h\}$. Therefore Z_h is the set of columns $(J_h(X), J_h(Y))$. For any set of columns $(J(X), J(Y))$, we represent the "row sum" for row i by $row^i(J(X), J(Y))$, which has the value $\sum_{j \in J(X)} X[i,j] + \sum_{j \in J(Y)} Y[i,j]$. We call Z_h an "column constrained table" because some rows i may have $row^i(J(X), J(Y)) \neq r_i$. However, all the column sums are satisfied by Z_h.

Step 1: We show we can modify Z_h by "deleting" at most d_m columns (including all of the X_j columns for $j \in R$) to bring the row sum for every row i below $r_i(1 - 1/n)$. We also show a dual result - if we "add" at most d_m columns to Z_h this brings the row sum for every row i above $r_i(1 + 1/n)$. Let $V_{i,h}$ and $W_{i,h}$ be defined as in (3.8) and (3.9).

First, instead of considering Z_h, consider $(\pi([n'] - V_{i,h})), \pi(V_{i,h}))$. This set of columns is the result of starting with the contingency table X, removing X_j for every reserved column $j \in R$, and then adding the w_j weights, for $j \in \pi(V_{i,h})$. By (3.8), we know that $|V_{i,h} \oplus \{1, \ldots, h\}| \leq 17m^2$ and

$$\begin{aligned}
row^i(\pi([n'] - V_{i,h}), \pi(V_{i,h})) &\leq (r_i - \sum_{j \in R} X[i,j]) + (h-1)\mu^i + 3c_m M_i \\
&= (r_i - \sum_{j \in R} X[i,j]) + 3c_m M_i \\
&\quad + \frac{h-1}{n'}(\sum_{j \in R} X[i,j] - Y[i,j]) \\
&= r_i - \frac{n' - h + 1}{n'}(\sum_{j \in R} X[i,j]) \\
&\quad - \frac{h-1}{n'}(\sum_{j \in R} Y[i,j]) + 3c_m M_i \\
&\leq r_i(1 - \frac{1}{n}) \qquad (a)
\end{aligned}$$

where the last step follows because (i) we know that $M_i \leq X[i,j]$ for every $j \in R_i^X$, and $R_i^X \subseteq R$, $|R_i^X| = 3c_m + 1$. Also $\max_{j \in R_i^X} X[i,j] \geq r_i/n$. Thus $3c_m M_i + r_i/n \leq \sum_{j \in R} X[i,j]$; (ii) similarly, $3c_m M_i +$

$r_i/n \leq \sum_{j \in R} Y[i,j]$; (iii) the convex combination $((n'-h+1)\sum_{j\in R} X[i,j] + (h-1)\sum_{j\in R} Y[i,j])/n'$ is at least $3c_m M_i + r_i/n$.

Now suppose we also "delete" $\pi\{1,\ldots,h\} \oplus \pi\{V_{i,h}\}$ from $(\pi([n'] - V_{i,h}), \pi(V_{i,h}))$.

Let $B_{i,h} = R \cup (\pi\{1,\ldots,h\} \oplus \pi\{V_{i,h}\})$.

Then

$$\begin{aligned} J_h(X) - B_{i,h} &= \pi\{h+1,\ldots,n'\} \cap \pi([n'] - V_{i,h}) \\ J_h(Y) - B_{i,h} &= \pi\{1,\ldots,h\} \cap \pi(V_{i,h}) \end{aligned}$$

By (a), we have $row^i(J_h(X) - B_{i,h}, J_h(Y) - B_{i,h}) \leq row^i(\pi([n'] - V_{i,h}), \pi(V_{i,h})) \leq r_i(1 - 1/n)$. Also, $|B_{i,h}| \leq 2m(3c_m + 1) + 1 + 17m^2$.

For the dual result, consider $(\pi([n'] - W_{i,h}) \cup R, \pi(W_{i,h}) \cup R)$. This set of columns is the result of starting with the contingency table X, adding the Y_j columns for $j \in R$, and then adding the w_j weights, for $j \in \pi(W_{i,h})$. By (3.9), we know that $|W_{i,h} \oplus \{1,\ldots,h\}| \leq 17m^2$ and

$$\begin{aligned} row^i(\pi([n'] - W_{i,h}) \cup R, \pi(W_{i,h}) \cup R) &\geq (r_i + \sum_{j\in R} Y[i,j]) + (h-1)\mu^i - 3c_m M_i \\ &= (r_i + \sum_{j\in R} Y[i,j]) - 3c_m M_i + \\ &\quad \frac{h-1}{n'}(\sum_{j\in R} X[i,j] - Y[i,j]) \\ &= r_i + \frac{n'-h+1}{n'}(\sum_{j\in R} Y[i,j]) + \\ &\quad \frac{h-1}{n'}(\sum_{j\in R} X[i,j]) - 3c_m M_i \\ &\geq r_i(1 + \frac{1}{n}) \qquad (b) \end{aligned}$$

where the last step follows because (i) $3c_m M_i + r_i/n \leq \sum_{j\in R} X[i,j]$; (ii) $3c_m M_i + r_i/n \leq \sum_{j\in R} Y[i,j]$; (iii) convex combination.

Suppose we add $\pi\{1,\ldots,h\} \oplus \pi\{W_{i,h}\}$ to $(\pi([n'] - W_{i,h}) \cup R, \pi(W_{i,h}) \cup R)$.

Let $C_{i,h} = R \cup (\pi\{1,\ldots,h\} \oplus \pi\{W_{i,h}\})$.

Then

$$\begin{aligned} J_h(X) \cup C_{i,h} &= \pi\{h+1,\ldots,n'\} \cup \pi([n'] - W_{i,h}) \cup R \\ J_h(Y) \cup C_{i,h} &= \pi\{1,\ldots,h\} \cup \pi(W_{i,h}) \cup R \end{aligned}$$

By (b), we have $row^i(J_h(X) \cup C_{i,h}, J_h(Y) \cup C_{i,h}) \geq row^i(\pi([n'] - W_{i,h}) \cup R, \pi(W_{i,h}) \cup R) \geq r_i(1 + 1/n)$. Also, $|C_{i,h}| \leq 2m(3c_m + 1) + 1 + 17m^2$.

Finally, define

$$D_h \quad =_{def} \quad (\cup_i B_{i,h}) \cup (\cup_i C_{i,h})$$

Consider Z_h with all of the columns in D_h "deleted". This is the table

$$Z_h^* \quad =_{def} \quad (J_h(X) - D_h, J_h(Y) - D_h)$$

Since $B_{i,h} \subseteq D_h$ for every i, we have $row^i(J_h(X) - D_h, J_h(Y) - D_h) \leq r_i(1 - 1/n)$ for all i. Also define

$$\bar{Z}_h^* \quad =_{def} \quad (J_h(X) \cup D_h, J_h(Y) \cup D_h)$$

Since $C_{i,h} \subseteq D_h$ for every i, we have $row^i(J_h(X) \cup D_h, J_h(Y) \cup D_h) \geq r_i(1 + 1/n)$ for all i.

Note that $|D_h| = d_m = 2m(3c_m + 1) + 1 + 34m^2$ (as defined previously). Also D_h contains all of R, including the index k.

Step 2: Now we show how to convert Z_h into a element of Ω. We focus on the "deleted" columns D_h, and show that by changing only the entries of the columns in D_h, we can obtain a contingency table $Z'_h \in \Sigma_{r,c}$. We will also show a dual result: that if we define $\bar{Z}_h$ to be the set of columns which contains X_j for every Y_j column in Z_h and contains Y_j for every X_j column in Z_h, we can show the same result for $\bar{Z}_h$ (we can construct a $\bar{Z}'_h$ in $\Sigma_{r,c}$ by changing d_m columns).

First let $\widehat{r}_i = row^i(J_h(X) - D_h, J_h(Y) - D_h)$, the partial row sum for row i of Z_h with the D_h columns removed. Define $s_i = r_i - \widehat{r}_i$ for all i, the sum for row i of the subtable that was removed from Z_h.

Note that $s_i \geq r_i/n$ for all i.

Let $N_h = \sum_{i=1}^{m} s_i = \sum_{j \in D_h} c_j$, by construction. We have two cases.

First suppose $N_h < 2(md_m)^2$. It is well-known that whenever the total of the row sums equals the total of the column sums, there is at least one contingency table satisfying these row and column sums (see Diaconis and Gangolli [5]). For this case we choose *any* set of modified values $Z'_h[i,j]$ for $j \in D_h$ such that $\sum_{i=1}^{m} Z'_h[i,j] = c_j$ for all $j \in D_h$ and $\sum_{j \in D_h} Z'_h[i,j] = s_i$ for all $1 \leq i \leq m$. Note that because $N_h < 2(md_m)^2$ we have $s_i < 2(md_m)^2$ for all i and therefore $r_i < 2n(md_m)^2$ for all i.

Alternatively, assume that $N_h \geq 2(md_m)^2$. As above, we are guaranteed that there is some set of $Z'_h[i,j]$ values for $j \in D_h$ that satisfy the row and column sums. But, for this case, we will need something stronger – we show we can modify the values of $Z_h[i,j]$ for the $j \in D_h$ columns in a *structured way* to obtain a subtable Z'_h satisfying the induced row sums s_i and the column sums c_j.

We already know that k is the index of the largest c_j for $j \in D_h$. Let ℓ be the index of the biggest s_i value. For every $i \neq \ell$ and every $j \in D_h - \{k\}$, we define $a_{i,j}$ in terms of the overall row sums and the column sums.

$$a_{i,j} =_{def} \lfloor \min\{r_i, c_j\}/n(d_m)^2 \rfloor$$

Since $Z_h[i,j]$ is either $X[i,j]$ or $Y[i,j]$, we know $Z_h[i,j] \leq \min\{r_i, c_j\}$. Therefore we can write

$$Z_h[i,j] = Q[i,j](a_{i,j} + 1) + R[i,j]$$

for $Q[i,j], R[i,j]$ non-negative integers, $Q[i,j] < n(d_m)^2$ and $0 \leq R[i,j] \leq a_{i,j}$ (unique), for every $i \neq \ell$ and every $j \in D_h - \{k\}$. We will show that by changing only the values of the $Q[i,j]$ to new values $Q'[i,j]$, we can obtain a subtable Z'_h satisfying the row sums s and the column sums.

Our analysis of the flow that we have constructed will rely on bounding the number of column constrained tables Z_h that can be transformed into the contingency table Z'_h (assuming we already know D_h, the d_m columns which have been modified to obtain Z'_h). If we know the subtable sum N_h is less than $2(md_m)^2$, then $r_i < 2n(md_m)^2$ for every i and therefore for every $j \in D_h$ and every row i, there are at most $2n(md_m)^2$ possible original values for $Z_h[i,j]$. In the general case when $N_h \geq 2(md_m)^2$, we do not have an upper bound on the original entries of the subtable. However, the method above for constructing the $Z'_h[i,j]$ values using $a_{i,j}$ gives an indirect upper bound for the number of Z_h that could be converted to Z'_h. For any true c.t. Z'_h and any selection of D_h columns which gives $N_h \geq 2(md_m)^2$, we can calculate $a_{i,j}$ for all $i \neq \ell$ and all $j \in D_h - \{k\}$. Then, for every row $i \neq \ell$ and every column j in $D_h - \{k\}$, we can find the original value of $R[i,j]$ using $R[i,j] = Z'_h[i,j] \bmod (a_{i,j} + 1)$. By definition of the $a_{i,j}$ values, there are at most $n(d_m)^2$ possible values for the original $Q[i,j]$, so there are only $(n(d_m)^2)^{(m-1)(d_m-1)}$ different ways of filling in the entire subtable on the $D_h - \{k\}$ columns.

We now return to our proof, since we still need to show that we can choose values $Q'[i,j]$ which ensure that the $Z'_h[i,j]$ values will satisfy row sums s_i and column sums c_j. Recall our assumption that $N_h \geq 2(md_m)^2$. It is well-known (see Dyer et al. [10]) that the row and column sums are satisfied by any integer matrix Z'_h which has $Z'_h[i,j] \geq 0$ for all i and also satisfies the following inequalities:

$$\sum_{i \neq \ell} Z'_h[i,j] \leq c_j \quad \text{for every } j \in D_h - \{k\} \tag{2}$$

$$\sum_{j \in D_h - \{k\}} Z'_h[i,j] \leq s_i \quad \text{for every } i \neq \ell \tag{3}$$

$$\sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} Z'_h[i,j] \geq N_h - s_\ell - c_k \tag{4}$$

Now define the $Q'[i,j]$ in terms of the induced row sums and the original column sums:

$$Q'[i,j] \quad =_{def} \quad \lfloor s_i c_j / N_h (a_{ij}+1) \rfloor$$

for all $i \neq \ell$ and all $j \in D_h - \{k\}$. Let $Z'_h[i,j] = Q'[i,j](a_{i,j}+1) + R[i,j]$. Note that $Q'[i,j] \geq 0$ for all i,j. We now prove that equalities 2, 3 and 4 are satisfied. Inequality (2):

$$\begin{aligned}
\sum_{i \neq \ell} Z'_h[i,j] &= \sum_{i \neq \ell} Q'[i,j](a_{i,j}+1) + \sum_{i \neq \ell} R[i,j] \\
&= \sum_{i \neq \ell} \lfloor s_i c_j / N_h (a_{ij}+1) \rfloor (a_{i,j}+1) + \sum_{i \neq \ell} R[i,j] \\
&\leq \sum_{i \neq \ell} s_i c_j / N_h + \sum_{i \neq \ell} R[i,j] \\
&\leq \sum_{i \neq \ell} s_i c_j / N_h + \sum_{i \neq \ell} a_{i,j} \\
&= c_j (1 - s_\ell / N_h) + \sum_{i \neq \ell} \lfloor \min\{r_i, c_j\} / n (d_m)^2 \rfloor \\
&\leq c_j (1 - s_\ell / N_h) + c_j (m-1) / n (d_m)^2 \\
&\leq c_j (1 - 1/m + 1/n d_m) \\
&\leq c_j.
\end{aligned}$$

The last two steps use the facts that (i) $s_\ell / N_h \geq 1/m$ (because s_ℓ is the largest row sum) and (ii) $d_m > m$.

Inequality (3): The early steps are similar to the proof for inequality (2). We get

$$\begin{aligned}
\sum_{j \in D_h - \{k\}} Z'_h[i,j] &\leq s_i (1 - c_k / N_h) + \sum_{j \in D_h - \{k\}} \lfloor \min\{r_i, c_j\} / n (d_m)^2 \rfloor \\
&\leq s_i (1 - c_k / N_h) + (d_m - 1) r_i / n (d_m)^2 \\
&\leq s_i (1 - 1/d_m) + s_i / d_m \\
&\leq s_i
\end{aligned}$$

where the second last step uses the fact that $c_k / N_h \geq 1/d_m$ (because c_k is the largest column sum in the subtable) and that $s_i \geq r_i / n$.

Inequality (4):

$$\begin{aligned}
\sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} Z'_h[i,j] &= \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} Q'[i,j](a_{i,j}+1) + \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} R[i,j] \\
&= \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} \lfloor s_i c_j / N_h (a_{i,j}+1) \rfloor (a_{i,j}+1) + \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} R[i,j] \\
&\geq \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} s_i c_j / N_h - \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} (a_{i,j}+1) + \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} 0 \\
&= (N_h - s_\ell)(N_h - c_k)/N_h - \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} (a_{i,j}+1) \\
&= (N_h - s_\ell - c_k) + s_\ell c_k / N_h - \sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} (a_{i,j}+1).
\end{aligned}$$

Next, note that

$$\begin{aligned}
s_\ell c_k / N_h &\geq N_h / (m d_m) \\
\sum_{i \neq \ell} \sum_{j \in D_h - \{k\}} (a_{i,j}+1) &\leq m d_m + m (N_h - c_k) / n (d_m)^2 \leq m d_m + N_h / (n d_m m),
\end{aligned}$$

by definition of the $a_{i,j}$ and by $d_m \geq m^2$. Therefore, to show that inequality (4) holds, it suffices to show

$$N_h/(md_m) \geq md_m + N_h/(nd_m m). \tag{5}$$

We note the following statements:

$$\begin{aligned} N_h/(md_m) &\geq md_m + N_h/(nd_m m) \\ \iff N_h(1-1/n)/(md_m) &\geq md_m \\ \impliedby N_h/2 &\geq (md_m)^2 \quad \text{(using that } n \geq 2) \\ \iff N_h &\geq 2(md_m)^2. \end{aligned}$$

This last statement holds by our previous assumption, therefore inequality (5) holds, and hence (4) is true, as required.

Therefore, in parallel with our path of column constrained tables, we have a path $X = Z'_0, Z'_1, \ldots, Z'_h, \ldots, Z'_{n'}$ such that Z'_h differs from Z_h in only d_m columns and $Z'_0, Z'_1, \ldots$ are true contingency tables. We can add another step to change $Z'_{n'}$ (using one step of the Markov chain) into Y. The amount of flow from X to Y that is routed along this path will be proportional to the probability that π is chosen.

Remember that the column constrained table Z_h contains exactly n columns and that $X_j \in Z_h \Leftrightarrow Y_j \notin Z_h$. Then if we define $\bar{Z}_h$ by the pair of sets $(J_h(Y), J_h(X))$, $\bar{Z}_h$ also contains exactly n columns and $X_j \in \bar{Z}_h \Leftrightarrow X_j \notin Z_h$. Now to calculate the row sums of $\bar{Z}_h$, consider $\bar{Z}_h$ with the columns of D_h removed. Then

$$\begin{aligned} row^i(J_h(Y) - D_h, J_h(X) - D_h) &= row^i(([n] - J_h(X)) - D_h), ([n] - J_h(Y)) - D_h) \\ &= row^i([n] - (J_h(X) \cup D_h), [n] - (J_h(Y) \cup D_h) \\ &= 2r_i - row^i(J_h(X) \cup D_h, J_h(Y) \cup D_h) \\ &\leq r_i(1 - 1/n) \end{aligned}$$

because we proved that $row^i(J_h(X) \cup D_h, J_h(Y) \cup D_h) \geq r_i(1 + 1/n)$ in Step 1.

Therefore, for $\bar{Z}_h$, there also exists a set of d_m columns (the same set of indices D_h) such that we can obtain a true contingency table $\bar{Z}'_h \in \Sigma_{r,c}$ by modifying these columns in the structured way described above.

Analysis: Next we show that the flow through any state $Z' \in \Omega$ is at most poly$(n)|\Omega|$. (Remark: We actually bound all of the flow *except* that due to pairs (X, Y) where $Z' = Y$. The total flow for all such pairs is only $|\Omega|$. Hence showing a bound of the form $poly(n)|\Omega|$ for the flow between remaining pairs where $Z' \neq Y$ provides us a similar bound for the total flow through Z'.) We assume Z' occurs as Z'_h for some pairs of contingency tables (X, Y) and some values of $\pi\{1, \ldots, h\}$. Again, we write $\bar{Z}_h$ for an column constrained table with X_j for $j \in J_h(Y)$ and Y_j for $j \in J_h(X)$. We claim that if we are given

(1) $\bar{Z}'_h \in \Omega$, a true contingency table obtained by changing d_m of the columns of $\bar{Z}_h$;

(2) the value of h and the set $\pi\{1, \ldots, h\}$ of columns already changed from X to Y in Z_h;

(3) The set D_h of d_m columns of Z_h which were modified to obtain Z'_h (note that the columns of $\bar{Z}_h$ that were modified are also the columns in D_h);

(4) The index ℓ of the largest induced row sum s_ℓ for the subtable of Z_h on D_h and the index ℓ' of the largest induced row sum for the subtable of $\bar{Z}_h$ on D_h;

(5) Two possibilities.
(i) If $N_h < 2(md_m)^2$, then we are given the original values of $Z_h[i, j]$ for all i, for all $j \in D_h$. Note we have $Z_h[i, j] \leq 2n(md_m)^2$;
(ii) Otherwise if $N_h \geq 2(md_m)^2$, we are given the original integers $Q[i, j]$ obtained from $Z_h[i, j]$ for every $i \neq \ell$ and every $j \in D_h - \{k\}$. Note we have $0 \leq Q[i, j] < n(d_m)^2$;

(5′) Two possibilities.
(i) If $\bar{N}_h < 2(md_m)^2$, then we are given the original values of $\bar{Z}_h[i, j]$ for all i, for all $j \in D_h$. Note we have $\bar{Z}_h[i, j] \leq 2n(md_m)^2$;
(ii) Otherwise if $\bar{N}_h \geq 2(md_m)^2$, we are given the original integers $\bar{Q}[i, j]$ obtained from $\bar{Z}_h[i, j]$ for every $i \neq \ell'$ and every $j \in D_h - \{k\}$. Note we have $0 \leq \bar{Q}[i, j] < n(d_m)^2$;

then we can construct X and Y. We refer to the information in (1)-(5′) as an "encoding" (for X and Y at the element $Z' \in \Omega$).

First concentrate on recovering Z_h. By (3) we know the submatrix D_h which has to be modified to obtain Z_h from Z'_h. Since we know the D_h columns we can calculate N_h. If $N_h < 2(md_m)^2$, then we have the original $Z_h[i,j]$ values for $j \in D_h$ and this gives us the entire idealized contingency table Z_h. If $N_h \geq 2(md_m)^2$ then we use (4) to identify ℓ. For every $i \neq \ell$ and $j \in D_h - \{k\}$ we calculate $a_{i,j}$ from r_i and c_j and we calculate $R[i,j] = Z'_h[i,j] \bmod (a_{i,j}+1)$. Finally, by (5) we have the original values of $Q[i,j]$ for all $i \neq \ell$ and all $j \in D_h - \{k\}$. Therefore we calculate $Z_h[i,j] = Q[i,j](a_{i,j}+1) + R[i,j]$. For columns $j \in D_h - \{k\}$, we can also calculate $Z_h[\ell,j]$ by subtracting the other values of $Z_h[i,j]$ (which we just calculated) from c_j. We still need to find the values $Z_h[i,k]$, but we defer this for the moment.

In a similar way, we use (3), (4) and (5′) to obtain all columns except column k of $\bar{Z}_h$ from $\bar{Z}'_h$ (given to us in (1)).

We now have all but column k of Z_h and $\bar{Z}_h$. Z_h contains column X_j for every $j \in J_h(X) = R \cup \pi\{h+1,\ldots,n'\}$ and contains column Y_j for every $j \in J_h(Y) = \pi\{1,\ldots,h\}$. $\bar{Z}_h$ contains column X_j for every $j \in J_h(Y) = [n] - J_h(X)$ and column Y_j for every $j \in J_h(X) = [n] - J_h(Y)$. By (2), we are given $\pi\{1,\ldots,h\}$. Then the c.t. X is the set of columns where

$$\begin{aligned} X_j &= \text{column } j \text{ of } Z_h \text{ for } j \in [n] - \pi\{1,\ldots,h\} \\ &= \text{column } j \text{ of } \bar{Z}_h \text{ for } j \in \pi\{1,\ldots,h\} \end{aligned}$$

The c.t. Y is the set of columns where

$$\begin{aligned} Y_j &= \text{column } j \text{ of } Z_h \text{ for } j \in \pi\{1,\ldots,h\} \\ &= \text{column } j \text{ of } \bar{Z}_h \text{ for } j \in [n] - \pi\{1,\ldots,h\} \end{aligned}$$

Thus for any $Z'_h \in \Omega$, we can construct all columns of X and Y except column k. Column k of each of these can then be recovered using the original r_i values. Thus, for any $Z'_h \in \Omega$, we can construct X and Y for any (X,Y) whose path passes through Z'_h, given the encoding (1)-(5′).

Now we bound the flow through our given Z'_h. Note that the flow through Z'_h is (bounded by) the number of possible choices for (1) and for (3)-(5′), times the amount of flow given by all possible choices for $\pi\{1,\ldots,h\}$ (given by (2)).

Since $\bar{Z}'_h \in \Omega$, there are $|\Omega|$ choices for (1).

There are $\binom{n}{d_m-1}$ possible D_h sets (choices for (3)), since D_h always contains k.

There are m^2 choices for (4).

Depending on the value of N_h, either there are at most $n(d_m)^2$ possible values for the original value of $Z_h[i,j]$ for $j \in D_h$ (obtained via $Q[i,j]$), or there are at most $2n(md_m)^2$ possible values for the original value of $Z_h[i,j]$ for $j \in D_h$. Therefore there are at most $(2n(md_m)^2)^{md_m}$ possible sets of $Z_h[i,j]$ for (5). The same upper bound holds for (5′).

The total number of choices for (4)-(5′) is $m^2(2n(md_m)^2)^{2md_m}$. We can write this as $C_m n^{2md_m}$, where C_m is the constant $2^{2m(d_m)}m^2(md_m)^{4md_m}$.

Recall that we are shipping one unit of flow from X to Y for all ordered pairs $(X,Y) \in \Omega^2$, and we want to find an upper bound on the amount of flow that passes through a given element $Z' \in \Omega$. The portion of flow from X to Y passing through Z' is determined by the distribution on the choices of permutations in (2). The permutation π defines the sequence $Z_0,\ldots,Z_n$ which, in turn, defines $Z'_0,\ldots,Z'_n$. We want to know how much flow passes through Z'_h for the choices (1),(3)-(5′) of the encoding. Therefore, we see the amount passing through Z'_h is bounded above by

$$|\Omega| C_m \binom{n}{d_m} n^{2md_m} \sum_h \sum_{U:\ |U|=h} \Pr[\pi\{1,\ldots,h\} = U]$$

where $\Pr[\pi\{1,...,h\} = U]$ is the probability that U is the set of the first h columns to be changed.

Therefore we need to bound $\sum_h \sum_{U:\ |U|=h} \Pr[\pi\{1,\ldots,h\} = U]$ for known Z_h, $\bar{Z}_h$ (following Morris). Note that since we have all of Z_h except column k and all of $\bar{Z}_h$ except column k, therefore for each value of $j \neq k$ we have $\{X_j, Y_j\}$. Also, since we know D_h we see there are at most 2^{d_m-1} possibilities for R (which

must include the unknown column k), and we shall sum over all of them. We then upper bound the flow coming from all permutations $\pi\{1,\ldots,h\}$ on the columns of $[n]-R$. There are at most $(2(n-|R|))^m \le (2n)^m$ possibilities for the vector M giving the M_i values and we like likewise sum over all such choices. For each choice of (possible) values $\{M_i\}$ we can compute $L_i = \{j : j \notin R, X[i,j] > M_i\} \cup \{j : j \notin R, Y[i,j] > M_i\}$. Note that this gives us S as well. By our remark on page 1, we know that $L_i \subseteq X$ or $L_i \subseteq Y$ for each i. Therefore there are only two possibilities for assigning all the L_i columns to X or Y (2^m choices for all of L).

Let $h_2 = h - |\pi\{1,\ldots,h\} \cap L|$. We bound the flow passing through Z'_h below. Note that the first summation is over all $(2(n-|R|))^m$ possibilities for M and all 2^m possible choices of L.

$$\begin{aligned}
& |\Omega| C_m \binom{n}{d_m} n^{2md_m} \sum_{R,M,L,h_2} \sum_{U \subseteq S:\ |U|=h_2} \Pr[\pi_2\{1,\ldots,h_2\} = U] \\
\le\ & |\Omega| C_m \binom{n}{d_m} n^{2md_m} 2^{d_m-1} (2n)^m 2^m n \sum_{U \subseteq S:\ |U|=h_2} \Pr[\pi_2\{1,\ldots,h_2\} = U] \\
\le\ & |\Omega| C_m \binom{n}{d_m} n^{2md_m} 2^{d_m-1} (2n)^m 2^m n \sum_{U \subseteq S:\ |U|=h_2} p_m(|S|) / \binom{|S|}{h_2} \\
\le\ & |\Omega| C_m \binom{n}{d_m} n^{2md_m} 2^{d_m-1} (2n)^m 2^m n \binom{|S|}{h_2} p_m(|S|) / \binom{|S|}{h_2} \\
\le\ & |\Omega| C_m \binom{n}{d_m} n^{2md_m} 2^{d_m-1} (2n)^m 2^m n p_m(|S|),
\end{aligned}$$

which is $poly(n)|\Omega|$.

Thus we know that the flow through any *state* is bounded by $\mathrm{poly}(n)|\Omega|$. In the application of Morris [17] this is already sufficient to prove polynomial time mixing, since the term $P(e)$ in the denominator of (1) is only polynomially small. However, for our heat-bath chain P_{HB}, it may be exponentially small, and further argument is required to establish rapid mixing.

To this end, let $e = (\omega, \omega')$ $(\omega, \omega' \in \Omega^2)$ be a (directed) transition of our heat-bath chain, with transition probability $P_{\mathrm{HB}}(e)$. Suppose that f_e units of flow are shipped along e in the multi-commodity flow defined above. We will disperse the flow through e by sending it from ω to ω' via a "random destination" ω''.

Let B be the set of columns on which ω and ω' disagree and let W be the set of all size $m \times (2d_m + 1)$ heat-bath windows which include B. Let Ω'' be the set of all contingency tables ω'' such that

1. There is a $U \in W$ which contains all the columns on which ω and ω'' differ, and
2. There is a $U' \in W$ which contains all the columns on which ω' and ω'' differ.

For each $\omega'' \in \Omega''$, we will route $f_e/|\Omega''|$ flow from ω to ω' via ω''. Note that this construction doubles the length of our flow paths, but no more.

The quantity shipped through (ω, ω'') from the original transition e in our multicommodity flow is $f_e/|\Omega''|$, which is at most $f_e n^{2d_m+1} P_{\mathrm{HB}}(\omega, \omega'')$. To see this, let K be the set of columns on which ω and ω'' differ. For every heat-bath window U, let $\Omega_\omega(U)$ denote the set of contingency tables which agree with ω, except possibly on window U and write

$$\begin{aligned}
P_{\mathrm{HB}}(\omega, \omega'') &= \sum_{U \supseteq K} \frac{1}{\binom{n}{2d_m+1}} \frac{1}{\Omega_\omega(U)} \\
&\ge \sum_{U \supseteq K, U \in W} \frac{1}{\binom{n}{2d_m+1}} \frac{1}{\Omega_\omega(U)} \\
&\ge \sum_{U \supseteq K, U \in W} \frac{1}{\binom{n}{2d_m+1}} \frac{1}{|\Omega''|} \\
&\ge \frac{1}{\binom{n}{2d_m+1}} \frac{1}{|\Omega''|},
\end{aligned}$$

where the last inequality follows from the fact that $\omega'' \in \Omega''$, so there is at least one U in the summation.

We call the transition (ω, ω'') a *type 1* transition and a transition (ω'', ω') a *type 2* transition.

We can now give an upper-bound for the total type 1 flow along the transition (ω, ω''). For each $e = (\omega, \omega')$, we ship at most $f_e n^{2d_m+1} P_{\mathrm{HB}}(\omega, \omega'')$ flow. Let f be the bound from above on the *total* flow that leaves node ω in our original multicommodity flow (so $f = \sum_e f_e$ where the sum is over transitions e which start at ω). Then the total amount now routed via ω'' is at most $f n^{2d_m+1} P_{\mathrm{HB}}(\omega, \omega'')$.

Using a symmetric argument, we can show that the total type 2 flow along the transition (ω'', ω') is at most $f n^{2d_m+1} P_{\mathrm{HB}}(\omega'', \omega')$.

Thus, the total flow in the new multicommodity flow along transition (ω, ω'') is at most $2 f n^{2d_m+1} P_{\mathrm{HB}}(\omega, \omega'')$. This is now sufficient for the right hand side of (1) to be polynomially bounded, since the (possibly small) $P_{\mathrm{HB}}(e)$ term cancels. This completes the proof of Theorem 8.

4 Mixing of the 2×2 chain

Theorem 8 shows that the Markov chain $\mathcal{M}_{\mathrm{HB}}$ is rapidly mixing. In this section we use the comparison method of Diaconis and Saloff-Coste [6] to show that the 2×2 chain $\mathcal{M}_{2\times 2}$ is also rapidly mixing. We briefly describe the comparison method, in the context of contingency tables, following the treatment of Vigoda [24]. For more details on the method, see [6], Randall and Tetali [21], and [24].

4.1 Setting up the comparison

Recall that P_{HB} denotes the transition matrix of the Markov chain $\mathcal{M}_{\mathrm{HB}}$ which was analyzed in Section 3. Let $E(P_{\mathrm{HB}})$ be the set of edges in the Markov kernel of the chain. That is, $E(P_{\mathrm{HB}}) = \{(X, Y) : P_{\mathrm{HB}}(X, Y) > 0\}$. Let $P_{2\times 2}$ denote the transition matrix of $\mathcal{M}_{2\times 2}$ and let $E(P_{2\times 2})$ denote the edge-set of its Markov kernel.

For every $(X, Y) \in E(P_{\mathrm{HB}})$ we define a set of paths $\Gamma_{X,Y}$ where each $\gamma \in \Gamma_{X,Y}$ is a path $X = \eta_0, \eta_1, \ldots, \eta_k = Y$, such that $(\eta_i, \eta_{i+1}) \in E(P_{2\times 2})$ for all $i \in \{0, \ldots, k-1\}$. Let $|\gamma|$ denote the length (number of edges) of the path. We also define a *flow* $f_{X,Y}$, which is a function from $\Gamma_{X,Y}$ to the positive reals satisfying the condition

$$\sum_{\gamma \in \Gamma_{X,Y}} f_{X,Y}(\gamma) = 1. \tag{6}$$

For each $(Z, Z') \in E(P_{2\times 2})$, define the quantity

$$A_{Z,Z'} = \frac{1}{P_{2\times 2}(Z, Z')} \sum_{(X,Y) \in E(P_{\mathrm{HB}})} \sum_{\substack{\gamma \in \Gamma_{X,Y} \text{ such} \\ \text{that } (Z,Z') \in \gamma}} |\gamma| f_{X,Y}(\gamma) P_{\mathrm{HB}}(X, Y).$$

We use the comparison theorem of Diaconis and Saloff-Coste, which says that

$$\tau_{\mathcal{M}_{2\times 2}}(\epsilon) \in O(\tau_{\mathcal{M}_{\mathrm{HB}}}(\epsilon) \log(|\Sigma_{r,c}|) \max_{(Z,Z') \in E(P_{2\times 2})} A_{Z,Z'}).$$

In our construction of the flow, we ensure that the length of each path $\gamma \in \Gamma_{X,Y}$ is bounded by a constant. Thus, the theorem of Diaconis and Saloff-Coste tells us that to establish rapid mixing, we need only define $f_{X,Y}$ for every $(X, Y) \in E(P_{\mathrm{HB}})$ such that Equation (6) is satisfied and also, for all $(Z, Z') \in E(P_{2\times 2})$, the following is satisfied:

$$\frac{1}{P_{2\times 2}(Z, Z')} \sum_{(X,Y) \in E(P_{\mathrm{HB}})} \sum_{\substack{\gamma \in \Gamma_{X,Y} \text{ such} \\ \text{that } (Z,Z') \in \gamma}} f_{X,Y}(\gamma) P_{\mathrm{HB}}(X, Y) \leq \mathrm{poly}(n). \tag{7}$$

It helps us to re-work Equation (7) before defining the flows. For $(X, Y) \in E(P_{\mathrm{HB}})$, let $\mathcal{W}(X, Y)$ be the set of all $m \times (2d_m + 1)$ "windows" such that X and Y agree outside of W, where a "window" is just a set of m rows and $2d_m + 1$ columns. Note that

$$P_{\mathrm{HB}}(X, Y) = \sum_{W \in \mathcal{W}(X,Y)} \frac{1}{\binom{n}{2d_m+1}} \frac{1}{|\Omega_X(W)|},$$

where $\Omega_X(W)$ is the set of all contingency tables that agree with X outside of W. We may view $P_{\mathrm{HB}}(X,Y)$ as an average of the quantities $1/|\Omega_X(W)|$ over all windows $W \in \mathcal{W}(X,Y)$. Therefore, we can pick some $W(X,Y) \in \mathcal{W}(X,Y)$ such that

$$P_{\mathrm{HB}}(X,Y) \leq \frac{1}{|\Omega_X(W(X,Y))|}. \tag{8}$$

The essential idea to keep in mind in what follows is that routing the unit flow $f_{X,Y}$ from X to Y is done using paths of contingency tables that differ from one another solely on (a part of) the chosen window $W(X,Y)$ satisfying (8).

For each $m \times (2d_m + 1)$ window W, let $E_W = \{(X,Y) \in P_{\mathrm{HB}} \mid W(X,Y) = W\}$. Later, when we define our flows, we do the following for every fixed window W: For every $(X,Y) \in E_W$, we define a flow $f_{X,Y}$ such that Equation (6) is satisfied. We also ensure that for all $(Z,Z') \in E(P_{2\times 2})$, the following is satisfied:

$$\frac{1}{P_{2\times 2}(Z,Z')} \sum_{(X,Y)\in E_W} \sum_{\substack{\gamma \in \Gamma_{X,Y} \text{ such} \\ \text{that } (Z,Z')\in\gamma}} f_{X,Y}(\gamma) P_{\mathrm{HB}}(X,Y) \leq \mathrm{poly}(n). \tag{9}$$

Since there are only polynomially-many windows W, equation (9) implies equation (7), and ensures rapid mixing.

For each window W, Section 4.2 shows how to define a flow $f^*_{X,Y}$ for every $(X,Y) \in E_W$ such that

$$\sum_{\gamma\in\Gamma_{X,Y}} f^*_{X,Y}(\gamma) = 1$$

and the total flow through any contingency table $Z \in \Sigma_{r,c}$ is in $O(|\Omega_X(W)|)$. We define $f_{X,Y}$ by modifying $f^*_{X,Y}$. Let $f^*_{X,Y}(Z)$ denote the amount of flow passing through the contingency table Z in the flow $f^*_{X,Y}$. Let $f^*(Z) = \sum_{(X,Y)\in E_W} f^*_{X,Y}(Z)$. Similarly, let $f_{X,Y}(Z,Z')$ denote the amount of flow passing through the transition (Z,Z') in the flow $f_{X,Y}$. Let $f(Z,Z') = \sum_{(X,Y)\in E_W} f_{X,Y}(Z,Z')$. Our construction of $f_{X,Y}$ from $f^*_{X,Y}$ ensures that for every $(Z,Z') \in E(P_{2\times 2})$, we have

$$f(Z,Z') \leq 2 f^*(Z) P_{2\times 2}(Z,Z') \binom{n}{2}\binom{m}{2}. \tag{10}$$

Thus, the left-hand-side of (9) is equal to

$$\frac{f(Z,Z')}{P_{2\times 2}(Z,Z')} P_{\mathrm{HB}}(X,Y) \leq \frac{f(Z,Z')}{P_{2\times 2}(Z,Z')} \frac{1}{|\Omega_X(W)|} \leq \frac{2 f^*(Z)\binom{n}{2}\binom{m}{2}}{|\Omega_X(W)|} \leq \mathrm{poly}(n),$$

where the first inequality comes from Equation (8), the second comes from Equation (10) which we establish shortly, and the third comes from the fact that $f^*(Z) \in O(|\Omega_X(W)|)$, which is established in Section 4.2. We will then have shown that Equation (9) is satisfied, as required, so the 2×2 heat bath chain is rapidly mixing on $\Sigma_{r,c}$.

We finish this section by showing how to construct $f_{X,Y}$ given the flow $f^*_{X,Y}$ and to establish (10). The method is similar to (but simpler than) the one that we used at the end of Section 3. For each $(Z,Z'') \in E(P_{2\times 2})$ (except $Z = Z''$) there is a *unique* 2×2 window $U(Z,Z'')$ on which Z and Z'' disagree. Let $\Omega_Z(U(Z,Z''))$ denote the set of all contingency tables that agree with Z (and Z'') everywhere off of the 2×2 window $U(Z,Z'')$. Let $f^*_{X,Y}(Z,Z'')$ be the flow that passes through (Z,Z'') in $f^*_{X,Y}$. For each $Z' \in \Omega_Z(U(Z,Z''))$, some of this flow is allocated to the path Z, Z', Z''. In particular, $f^*_{X,Y}(Z,Z'')/|\Omega_Z(U(Z,Z''))|$ flow is sent this way.

Now $f_{X,Y}(Z,Z') \leq 2\sum_{Z''\in U(Z,Z')} \frac{f^*_{X,Y}(Z,Z'')}{|\Omega_Z(U(Z,Z''))|}$, where the first "2" comes from the fact that we must consider paths Z, Z', Z'' above, and also paths that end in edge Z, Z'. The right-hand side is at most

$$2\frac{f^*_{X,Y}(Z)}{|\Omega_Z(U(Z,Z'))|} \leq 2 f^*_{X,Y}(Z) P_{2\times 2}(Z,Z') \binom{n}{2}\binom{m}{2},$$

so (10) holds.

By considering all $m \times (2d_m + 1)$ sized windows which contain the two columns on which Z and Z' differ, we can see that for each $(Z, Z') \in E(P_{2\times 2})$, we have

$$A_{Z,Z'} \leq C \binom{n}{2} \binom{n-2}{2d_m - 1}$$

where the constant C accounts for the maximum length of any (X, Y) path for $(X, Y) \in E(P_{\mathrm{HB}})$, and the constant factors arising in the bound for the flow $f^*(Z)$ over a single $m \times (2d_m + 1)$ window W. Therefore, we have the following theorem:

Theorem 9

$$\tau_{\mathcal{M}_{2\times 2}}(\epsilon) \in O(\tau_{\mathcal{M}_{\mathrm{HB}}}(\epsilon) \log(|\Sigma_{r,c}|) n^{2d_m+1}).$$

4.2 Defining $f^*(X, Y)$

In this section we define a flow $f^*_{X,Y}$ for every $(X, Y) \in E_W$ such that $\sum_{\gamma \in \Gamma_{X,Y}} f^*_{X,Y}(\gamma) = 1$ and the total flow through any contingency table Z, due to pairs in E_W, is in $O(|\Omega_X(W)|)$.

Throughout the entire section, we therefore focus on some fixed $m \times (2d_m + 1)$ sized window W of the larger $m \times n$ table. Without loss of generality (and to make our notation simpler in what follows), we assume that W includes the first $2d_m + 1$ columns of the table. This window W has induced row sums ρ_i (for $i \in [m]$) and induced column sums ζ_j (for $j \in [2d_m + 1]$). For convenience we also set $\delta = 2d_m + 1$.

Let $\rho = (\rho_1, \ldots, \rho_m)$, $\zeta = (\zeta_1, \ldots, \zeta_\delta)$ be the lists of induced row and column sums. Let $\Sigma_{\rho,\zeta}$ denote the set of $m \times \delta$ contingency tables with rows sums ρ and column sums ζ, and let N_W denote the table sum. Let $\Upsilon, \Psi \in \Sigma_{\rho,\zeta}$. We show how to route a unit of flow between Υ and Ψ using a path of contingency tables that differ by 2×2 heat bath moves. This flow lifts in the obvious fashion to transitions $(X, Y) \in E(P_{\mathrm{HB}})$, giving us the flow $f^*_{X,Y}$ required in the previous section. In other words, we simply use the exact same sequence of 2×2 transitions on the window $W(X, Y)$, keeping everything outside this window fixed (where X and Y agree anyway).

If $N_W < (2m\delta)^2$ then $|\Sigma_{\rho,\zeta}| \in O(1)$, so it doesn't really matter how we route flow between Υ and Ψ. For example, it suffices to fix each square in the contingency table in lexicographic order. Each path in the resulting flow is of length $O(1)$ and there are $O(1)$ pairs (Υ, Ψ) of contingency tables, so the desired bound is easily established. This is similar to, but simpler than, what we do later in Section 4.2.3. Thus, from now on we assume $N_W \geq (2m\delta)^2$ and we show how to construct a flow between Υ and Ψ in $\Sigma_{\rho,\zeta}$.

Without loss of generality we may assume that the row totals are sorted into non-descending order and that the column totals are also sorted into non-descending order. Therefore ρ_m is the largest row sum and ζ_δ is the largest column sum.

As we did in section 3, we view the space $\Sigma_{\rho,\zeta}$ of contingency tables as the $(m-1)(\delta-1)$-dimensional space of integer matrices Φ that satisfy $\Phi[i, j] \geq 0$ for all $i \in [m-1]$ and all $j \in [\delta - 1]$ and also satisfy the following inequalities (see Dyer, Kannan and Mount [10]):

$$\sum_{i=1}^{m-1} \Phi[i, j] \leq \zeta_j \quad \text{for every } j \in [\delta - 1] \tag{11}$$

$$\sum_{j=1}^{\delta-1} \Phi[i, j] \leq \rho_i \quad \text{for every } i \in [m - 1] \tag{12}$$

$$\sum_{i=1}^{m-1} \sum_{j=1}^{\delta-1} \Phi[i, j] \geq N_W - \rho_m - \zeta_\delta. \tag{13}$$

Let

$$\alpha_{i,j} = \left\lfloor \frac{\min\{\rho_i, \zeta_j\}}{m^2\delta^2} \right\rfloor \tag{14}$$

for all $i \in [m-1]$, $j \in [\delta - 1]$.

Note that $\max_{j\in[\delta-1]} \alpha_{i,j} = \alpha_{i,\delta-1}$ for all i; that $\max_{i\in[m-1]} \alpha_{i,j} = \alpha_{m-1,j}$ for all j; and that $\max_{i\in[m-1],j\in[\delta-1]} \alpha_{i,j} = \alpha_{m-1,\delta-1}$.

For any contingency table $\Phi \in \Sigma_{\rho,\zeta}$, and any $i \in [m-1]$, $j \in [\delta-1]$, we can write

$$\Phi[i,j] = Q[i,j](\alpha_{i,j}+1) + R[i,j],$$

for a unique integer $R[i,j]$ satisfying $0 \le R[i,j] \le \alpha_{i,j}$, and a unique integer $Q[i,j]$.

Let

$$Q^*[i,j] \quad = \quad \left\lfloor \frac{\rho_i \zeta_j}{N_W(\alpha_{i,j}+1)} \right\rfloor \tag{15}$$

for all $i \in [m-1]$, $j \in [\delta-1]$.

If Φ is a contingency table such that $Q[i,j] = Q^*[i,j]$ for every $i \in [m-1]$, $j \in [\delta-1]$, then we say that Φ belongs to the *inner domain* of $\Sigma_{\rho,\zeta}$.

Let $\Upsilon, \Psi \in \Sigma_{\rho,\zeta}$. We are almost ready to start building a path from Υ to Ψ that uses 2×2 heat bath transitions. We use the following lemma.

Lemma 10 *Let $\Phi^*[i,j]$ be defined by $\Phi^*[i,j] = Q^*[i,j](\alpha_{i,j}+1) + R[i,j]$, for any non-negative integers $R[i,j] \le \alpha_{i,j}$, for all $i \in [m-1]$, $j \in [\delta-1]$. Then*

(i) $\Phi^* \in \Sigma_{\rho,\zeta}$.

(ii) Under the assumption that $N_W \ge (2m\delta)^2$, we have the following:

$$\begin{aligned}
\Phi^*[i,\delta] &\ge (\alpha_{i,\delta-1}+1) \text{ for all } i \in [m-1] \\
\Phi^*[m,j] &\ge (\alpha_{m-1,j}+1) \text{ for all } j \in [\delta-1] \\
\Phi^*[m,\delta] &\ge 2(\alpha_{m-1,\delta-1}+1)
\end{aligned}$$

Proof: (i) This is the same as the proof of Equations (2), (3), and (4).

(ii) By definition,

$$\begin{aligned}
\Phi^*[i,\delta] &= \rho_i - \sum_{j=1}^{\delta-1} \Phi^*[i,j] \\
&= \rho_i - \sum_{j=1}^{\delta-1} \left(\left\lfloor \frac{\rho_i \zeta_j}{N_W(\alpha_{i,j}+1)} \right\rfloor (\alpha_{i,j}+1) + R[i,j] \right) \\
&\ge \rho_i - \sum_{j=1}^{\delta-1} \left(\frac{\rho_i \zeta_j}{N_W} + \alpha_{i,j} \right) \\
&= \frac{\rho_i \zeta_\delta}{N_W} - \sum_{j=1}^{\delta-1} \alpha_{i,j} \\
&\ge \frac{\rho_i \zeta_\delta}{N_W} - \frac{\delta \rho_i}{m^2\delta^2} \\
&\ge \frac{\rho_i}{\delta} - \frac{\rho_i}{m^2\delta} \\
&\ge \frac{\rho_i}{m\delta} \\
&\ge 2\alpha_{i,\delta-1}.
\end{aligned}$$

Then if $\alpha_{i,\delta-1} \ge 1$, we automatically have $\Phi^*[i,\delta] \ge (\alpha_{i,\delta-1}+1)$. However if $\alpha_{i,\delta-1} = 0$, then $\alpha_{i,j} = 0$ for all $j \in [\delta-1]$. There are two subcases. The first subcase is when $\rho_i\zeta_\delta < N_W$. Then $\rho_i\zeta_j < N_W$ for all $j \in [\delta-1]$, and $\Phi^*[i,j] = 0$ for all $j \in [\delta-1]$. Then $\Phi^*[i,\delta] = \rho_i \ge 1$. If $\rho_i\zeta_\delta \ge N_W$, then the fourth line of the derivation above (with $\alpha_{i,j} = 0$) gives $\Phi^*[i,\delta] \ge \rho_i\zeta_\delta/N_W \ge 1$. So in either subcase we have $\Phi^*[i,\delta] \ge 1 = (\alpha_{i,\delta-1}+1)$.

Using a similar argument we get

$$\Phi^*[m,j] \quad \geq \quad (\alpha_{m-1,j}+1).$$

By definition

$$\begin{aligned}
\Phi^*[m,\delta] &= \zeta_\delta - \sum_{i=1}^{m-1} \Phi^*[i,\delta] \\
&= \zeta_\delta - \sum_{i=1}^{m-1} \left(\rho_i - \sum_{j=1}^{\delta-1} \Phi^*[i,j] \right) \\
&= (\rho_m + \zeta_\delta - N_W) + \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} \left(\left\lfloor \frac{\rho_i \zeta_j}{N_W(\alpha_{i,j}+1)} \right\rfloor (\alpha_{i,j}+1) + R[i,j] \right) \\
&\geq (\rho_m + \zeta_\delta - N_W) + \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} \left(\left\lfloor \frac{\rho_i \zeta_j}{N_W(\alpha_{i,j}+1)} \right\rfloor (\alpha_{i,j}+1) \right) \\
&\geq (\rho_m + \zeta_\delta - N_W) + \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} \left(\frac{\rho_i \zeta_j}{N_W} - (\alpha_{i,j}+1) \right) \\
&= \frac{\rho_m \zeta_\delta}{N_W} - \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} (\alpha_{i,j}+1) \\
&\geq \frac{\rho_m \zeta_\delta}{N_W} - \frac{m(N_W - \zeta_\delta)}{m^2\delta^2} - (m-1)(\delta-1) \\
&\geq \frac{N_W}{m\delta} - \frac{N_W}{m\delta^2} - (m-1)(\delta-1) \\
&\geq \frac{N_W}{4m\delta} + (\frac{N_W}{2m\delta} - \frac{N_W}{m\delta^2}) + (\frac{N_W}{4m\delta} - (m-1)(\delta-1)) \\
&\geq \frac{2\zeta_{\delta-1}}{4m\delta} + 0 + (\delta + m - 1) \\
&\geq 2\alpha_{m-1,\delta-1} + 2,
\end{aligned}$$

as required. □

For every $i \in [m-1]$, $j \in [\delta-1]$, write

$$\Upsilon[i,j] = Q_\Upsilon[i,j](\alpha_{i,j}+1) + R_\Upsilon[i,j],$$

for the unique integer $R_\Upsilon[i,j]$ that also satisfies $0 \leq R_\Upsilon[i,j] \leq \alpha_{i,j}$.

Similarly, for every $i \in [m-1]$, $j \in [\delta-1]$, write

$$\Psi[i,j] = Q_\Psi[i,j](\alpha_{i,j}+1) + R_\Psi[i,j],$$

for the unique $R_\Psi[i,j]$ that also satisfies $0 \leq R_\Psi[i,j] \leq \alpha_{i,j}$.

The routing from Υ to Ψ proceeds in two stages. In the first phase of the routing, we route flow from Υ to the table Υ^* in the inner domain of $\Sigma_{\rho,\zeta}$ such that $\Upsilon^*[i,j] = Q^*[i,j](\alpha_{i,j}+1) + R_\Upsilon[i,j]$ holds for every $i \in [m-1]$, $j \in [\delta-1]$. Note $\Upsilon^* \in \Sigma_{\rho,\zeta}$ using the previous lemma.

By defining a similar path between Ψ and Ψ^* and then reversing all the edges, we can route flow from some Ψ^* in the inner domain of $\Sigma_{\rho,\zeta}$ to Ψ such that $\Psi^*[i,j] = Q^*[i,j](\alpha_{i,j}+1) + R_\Psi[i,j]$ holds for every $i \in [m-1]$, $j \in [\delta-1]$. Similarly, we also have $\Psi^* \in \Sigma_{\rho,\zeta}$.

In the second phase of the routing we will route flow from Υ^* to Ψ^* by changing the $R_\Upsilon[i,j]$ to the $R_\Psi[i,j]$ values.

4.2.1 Phase 1

We show how to route Υ to Υ^*, by only changing $Q[i,j]$ values at any step. We define the following metric on pairs $\Phi, \Phi' \in \Sigma_{\rho,\zeta}$:

$$\begin{aligned}
d(\Phi,\Phi') &= d_1(\Phi,\Phi') + d_2(\Phi,\Phi') + d_3(\Phi,\Phi') + d_4(\Phi,\Phi') \qquad \text{where} \\
d_1(\Phi,\Phi') &= \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} |\Phi[i,j] - \Phi'[i,j]| \\
d_2(\Phi,\Phi') &= \frac{N_W+1}{3N_W}\left(\sum_{i=1}^{m-1} |\Phi[i,\delta] - \Phi'[i,\delta]|\right) \\
d_3(\Phi,\Phi') &= \frac{N_W+1}{3N_W}\left(\sum_{j=1}^{\delta-1} |\Phi[m,j] - \Phi'[m,j]|\right) \\
d_4(\Phi,\Phi') &= \frac{N_W+1}{3N_W}|\Phi[m,\delta] - \Phi'[m,\delta]|
\end{aligned}$$

This metric is used to show the path we construct is moving us closer to Υ^*.

We route Υ to Υ^* as a series of moves. Let Υ' be any interim contingency table on the path from Υ to Υ^*. We choose our next move from one of four cases.

Case a: Suppose that $\Upsilon'[m,\delta] < (\alpha_{m-1,\delta-1}+1)$. Then we perform a move to make $\Upsilon'[m,\delta]$ bigger (because we need to "leave room" for our other cases). In the full version, we show that there is at least one $\ell \in [m-1]$ such that $\Upsilon'[\ell,\delta] \geq \Upsilon^*[\ell,\delta] + (\alpha_{\ell,\delta-1}+1)$.

Note that by lemma 10, in this case $\Upsilon'[m,\delta] < \Upsilon^*[m,\delta] - (\alpha_{m-1,\delta-1}+1)$. Suppose (for contradiction) that there is no ℓ as described above, so that

$$\Upsilon'[i,\delta] \quad < \quad \Upsilon^*[i,\delta] + (\alpha_{i,\delta-1}+1) \;\text{ for } i \in [m-1].$$

By definition

$$\begin{aligned}
\Upsilon^*[i,\delta] &= \rho_i - (\sum_{j=1}^{\delta-1} \Upsilon^*[i,j]) \\
&= \rho_i - \left(\sum_{j=1}^{\delta-1}\left(\left\lfloor \frac{\rho_i\zeta_j}{N_W(\alpha_{i,j}+1)}\right\rfloor (\alpha_{i,j}+1) + R_\Upsilon[i,j]\right)\right).
\end{aligned}$$

Now

$$\begin{aligned}
\Upsilon'[m,\delta] &= \zeta_\delta - (\sum_{i=1}^{m-1} \Upsilon'[i,\delta]) \\
&> \zeta_\delta - \sum_{i=1}^{m-1}\left(\rho_i - \left(\sum_{j=1}^{\delta-1}\Upsilon^*[i,j]\right) + (\alpha_{i,\delta-1}+1)\right) \\
&= (\zeta_\delta + \rho_m - N_W) + \sum_{i=1}^{m-1}\left(\left(\sum_{j=1}^{\delta-1}\Upsilon^*[i,j]\right) - (\alpha_{i,\delta-1}+1)\right). \qquad (16)
\end{aligned}$$

Expanding $\sum_{i=1}^{m-1}((\sum_{j=1}^{\delta-1}\Upsilon^*[i,j]) - (\alpha_{i,\delta-1}+1))$

$$
\begin{aligned}
&= \sum_{i=1}^{m-1}\left(\sum_{j=1}^{\delta-1}\left(\left\lfloor\frac{\rho_i\zeta_j}{N_W(\alpha_{i,j}+1)}\right\rfloor(\alpha_{i,j}+1)+R_\Upsilon[i,j]\right)-(\alpha_{i,\delta-1}+1)\right)\\
&\geq \sum_{i=1}^{m-1}\left(\sum_{j=1}^{\delta-1}\left(\frac{\rho_i\zeta_j}{N_W}-(\alpha_{i,j}+1)\right)-(\alpha_{i,\delta-1}+1)\right)\\
&= (N_W-\rho_m-\zeta_\delta)+\frac{\rho_m\zeta_\delta}{N_W}-\sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1}(\alpha_{i,j}+1)-\sum_{i=1}^{m-1}(\alpha_{i,\delta-1}+1)\\
&\geq (N_W-\rho_m-\zeta_\delta)+\frac{\rho_m\zeta_\delta}{N_W}-\frac{m(N_W-\zeta_\delta)}{m^2\delta^2}-(m-1)(\delta-1)-\frac{m\zeta_{\delta-1}}{m^2\delta^2}-(m-1)\\
&\geq (N_W-\rho_m-\zeta_\delta)+\frac{N_W}{m\delta}-\frac{N_W}{m\delta^2}-m\delta+\delta\\
&\geq (N_W-\rho_m-\zeta_\delta)+\frac{N_W}{2m\delta}-m\delta+\delta\\
&\geq (N_W-\rho_m-\zeta_\delta)+\frac{N_W}{4m\delta}+\delta\\
&\geq (N_W-\rho_m-\zeta_\delta)+(\alpha_{m-1,\delta-1}+1) \qquad (17)
\end{aligned}
$$

where the second last line follows by $N_W \geq (2m\delta)^2$, and the last line follows from the definition of $\alpha_{i,j}$ and by $\delta \geq m \geq 2$.

Combining (16) and (17), we have a contradiction to our original assumption that $\Upsilon'[m,\delta] < (\alpha_{m-1,\delta-1}+1)$. Therefore it must be that $\Upsilon'[\ell,\delta] \geq \Upsilon^*[\ell,\delta]+(\alpha_{\ell,\delta-1}+1)$ for some $\ell \in [m-1]$.

Similarly, there must be some $k \in [\delta-1]$ such that $\Upsilon'[m,k] \geq \Upsilon^*[m,k]+(\alpha_{m-1,k}+1)$.

In this case we add the transition $(\Upsilon' \to \Upsilon'')$ to our path from Υ to Υ^*, where Υ'' is identical to Υ' except for the following entries:

$$
\begin{aligned}
\Upsilon''[\ell,k] &= \Upsilon'[\ell,k]+(\alpha_{\ell,k}+1) \quad & \Upsilon''[\ell,\delta] &= \Upsilon'[\ell,\delta]-(\alpha_{\ell,k}+1)\\
\Upsilon''[m,k] &= \Upsilon'[m,k]-(\alpha_{\ell,k}+1) \quad & \Upsilon''[m,\delta] &= \Upsilon'[m,\delta]+(\alpha_{\ell,k}+1)
\end{aligned}
$$

Now we show that $d(\Upsilon'',\Upsilon^*) < d(\Upsilon',\Upsilon^*)$.

$$
\begin{aligned}
d_1(\Upsilon'',\Upsilon^*) &= \textstyle\sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1}|\Upsilon''[i,j]-\Upsilon^*[i,j]| &\leq\ & d_1(\Upsilon',\Upsilon^*)+(\alpha_{\ell,k}+1)\\
d_2(\Upsilon'',\Upsilon^*) &= \tfrac{N_W+1}{3N_W}\left(\textstyle\sum_{i=1}^{m-1}|\Upsilon''[i,\delta]-\Upsilon^*[i,\delta]|\right) &=\ & d_2(\Upsilon',\Upsilon^*)-\tfrac{N_W+1}{3N_W}(\alpha_{\ell,k}+1)\\
d_3(\Upsilon'',\Upsilon^*) &= \tfrac{N_W+1}{3N_W}\left(\textstyle\sum_{j=1}^{\delta-1}|\Upsilon''[m,j]-\Upsilon^*[m,j]|\right) &=\ & d_3(\Upsilon',\Upsilon^*)-\tfrac{N_W+1}{3N_W}(\alpha_{\ell,k}+1)\\
d_4(\Upsilon'',\Upsilon^*) &= \tfrac{N_W+1}{3N_W}|\Upsilon''[m,\delta]-\Upsilon^*[m,\delta]| &=\ & d_4(\Upsilon',\Upsilon^*)-\tfrac{N_W+1}{3N_W}(\alpha_{\ell,k}+1)
\end{aligned}
$$

where the equation for d_4 follows by lemma 10.

So $d(\Upsilon'',\Upsilon^*) \leq d(\Upsilon',\Upsilon^*)+(\alpha_{\ell,k}+1)-(\alpha_{\ell,k}+1)(N_W+1)/N_W \leq d(\Upsilon',\Upsilon^*)-(\alpha_{\ell,k}+1)/N_W < d(\Upsilon',\Upsilon^*)$.

Case b: We execute (b) whenever Case (a) does not hold and when *either* $\Upsilon'[\ell,\delta] < \Upsilon^*[\ell,\delta]$ for some $\ell \in [m-1]$ *or* $\Upsilon'[m,j] < \Upsilon^*[m,j]$ for some $j \in [\delta-1]$. Assume the former without loss of generality. If this is the case, then there must be at least one $k \in [\delta-1]$ such that $\Upsilon'[\ell,k] > \Upsilon^*[\ell,k]$. Since we only change $Q[i,j]$ values during our routing, we know that $\Upsilon'[\ell,k] \geq \Upsilon^*[\ell,k]+(\alpha_{\ell,k}+1)$. Also, since we are not in Case (a), we know $\Upsilon'[m,\delta] \geq (\alpha_{m-1,\delta-1}+1) \geq (\alpha_{\ell,k}+1)$.

In this case we add the transition $(\Upsilon' \to \Upsilon'')$ in our path from Υ to Υ^*, where Υ'' is identical to Υ' except for the following entries:

$$
\begin{aligned}
\Upsilon''[\ell,k] &= \Upsilon'[\ell,k]-(\alpha_{\ell,k}+1) \quad & \Upsilon''[\ell,\delta] &= \Upsilon'[\ell,\delta]+(\alpha_{\ell,k}+1)\\
\Upsilon''[m,k] &= \Upsilon'[m,k]+(\alpha_{\ell,k}+1) \quad & \Upsilon''[m,\delta] &= \Upsilon'[m,\delta]-(\alpha_{\ell,k}+1)
\end{aligned}
$$

Now we show $d(\Upsilon'', \Upsilon^*) < d(\Upsilon', \Upsilon^*)$.

$$\begin{array}{lcl} d_1(\Upsilon'', \Upsilon^*) = \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} |\Upsilon''[i,j] - \Upsilon^*[i,j]| = d_1(\Upsilon', \Upsilon^*) - (\alpha_{\ell,k}+1) \\ d_2(\Upsilon'', \Upsilon^*) = \frac{N_W+1}{3N_W}\left(\sum_{i=1}^{m-1} |\Upsilon''[i,\delta] - \Upsilon^*[i,\delta]|\right) \leq d_2(\Upsilon', \Upsilon^*) + \frac{N_W+1}{3N_W}(\alpha_{\ell,k}-1) \\ d_3(\Upsilon'', \Upsilon^*) = \frac{N_W+1}{3N_W}\left(\sum_{j=1}^{\delta-1} |\Upsilon''[m,j] - \Upsilon^*[m,j]|\right) \leq d_3(\Upsilon', \Upsilon^*) + \frac{N_W+1}{3N_W}(\alpha_{\ell,k}+1) \\ d_4(\Upsilon'', \Upsilon^*) = \frac{N_W+1}{3N_W}|\Upsilon''[m,\delta] - \Upsilon^*[m,\delta]| \leq d_4(\Upsilon', \Upsilon^*) + \frac{N_W+1}{3N_W}(\alpha_{\ell,k}+1) \end{array}$$

where the expression for $d_2(\Upsilon'', \Upsilon^*)$ follows by $\Upsilon'[\ell,\delta] \leq \Upsilon^*[\ell,\delta] - 1$.

Then $d(\Upsilon'', \Upsilon^*) \leq d(\Upsilon', \Upsilon^*) - (\alpha_{\ell,k}+1) + (\alpha_{\ell,k}+1)(N_W+1)/N_W - 2(N_W+1)/3N_W$. This is at most $d(\Upsilon', \Upsilon^*) + (\alpha_{\ell,k}+1)/N_W - 2(N_W+1)/3N_W$. Then because $\alpha_{\ell,k} \leq N_W/(\delta m)^2$, we have $d(\Upsilon'', \Upsilon^*) \leq d(\Upsilon', \Upsilon^*) + 1/(m\delta)^2 + 1/3N_W - 2/3$, and using $N_W \geq (2\delta m)^2$, we obtain $d(\Upsilon'', \Upsilon^*) < d(\Upsilon', \Upsilon^*)$.

Case c: We execute (c) when cases (a)-(b) do not hold and *either* $\Upsilon'[\ell,\delta] > \Upsilon^*[\ell,\delta]$ for some $\ell \in [m-1]$ *or* $\Upsilon'[m,j] > \Upsilon^*[m,j]$ for some $j \in [\delta-1]$. Assume the former without loss of generality. Then there must exist some $k \in [\delta-1]$ such that $\Upsilon'[\ell,k] < \Upsilon^*[\ell,k]$, and since we can only change the $\Upsilon[\ell,k]$ values by factors of $(\alpha_{\ell,k}+1)$, we have $\Upsilon'[\ell,k] \leq \Upsilon^*[\ell,k] + (\alpha_{\ell,k}+1)$. Note that by $\Upsilon'[\ell,\delta] > \Upsilon^*[\ell,\delta]$ and by part (ii) of lemma 10, we know $\Upsilon'[\ell,\delta] \geq (\alpha_{\ell,\delta-1}+1) \geq (\alpha_{\ell,k}+1)$. Because case (b) does not hold, we know $\Upsilon'[m,k] \geq \Upsilon^*[m,k]$, and by part (ii) of lemma 10, this is at least $(\alpha_{m-1,k}+1) \geq (\alpha_{\ell,k}+1)$.

Therefore we can perform the transition $(\Upsilon' \to \Upsilon'')$, where Υ'' is identical to Υ' except for the following entries:

$$\begin{array}{ll} \Upsilon''[\ell,k] = \Upsilon'[\ell,k] + (\alpha_{\ell,k}+1) & \Upsilon''[\ell,\delta] = \Upsilon'[\ell,\delta] - (\alpha_{\ell,k}+1) \\ \Upsilon''[m,k] = \Upsilon'[m,k] - (\alpha_{\ell,k}+1) & \Upsilon''[m,\delta] = \Upsilon'[m,\delta] + (\alpha_{\ell,k}+1) \end{array}$$

Now we show $d(\Upsilon'', \Upsilon^*) < d(\Upsilon', \Upsilon^*)$.

$$\begin{array}{lcl} d_1(\Upsilon'', \Upsilon^*) = \sum_{i=1}^{m-1}\sum_{j=1}^{\delta-1} |\Upsilon''[i,j] - \Upsilon^*[i,j]| = d_1(\Upsilon', \Upsilon^*) - (\alpha_{\ell,k}+1) \\ d_2(\Upsilon'', \Upsilon^*) = \frac{N_W+1}{3N_W}\sum_{i=1}^{m-1} |\Upsilon''[i,\delta] - \Upsilon^*[i,\delta]| \leq d_2(\Upsilon', \Upsilon^*) + \frac{N_W+1}{3N_W}(\alpha_{\ell,k}-1) \\ d_3(\Upsilon'', \Upsilon^*) = \frac{N_W+1}{3N_W}\sum_{j=1}^{\delta-1} |\Upsilon''[m,j] - \Upsilon^*[m,j]| \leq d_3(\Upsilon', \Upsilon^*) + \frac{N_W+1}{3N_W}(\alpha_{\ell,k}+1) \\ d_4(\Upsilon'', \Upsilon^*) = \frac{N_W+1}{3N_W}|\Upsilon''[m,\delta] - \Upsilon^*[m,\delta]| \leq d_4(\Upsilon', \Upsilon^*) + \frac{N_W+1}{3N_W}(\alpha_{\ell,k}+1) \end{array}$$

where the bound on $d_2(\Upsilon'', \Upsilon^*)$ follows by $\Upsilon'[\ell,\delta] \geq \Upsilon^*[\ell,\delta] + 1$.

Case d: This is the case when $\Upsilon'[i,\delta] = \Upsilon^*[i,\delta]$ for all $i \in [m-1]$ and $\Upsilon'[m,j] = \Upsilon^*[m,j]$ for all $j \in [\delta-1]$ (so neither of cases (b) or (c) hold), but $\Upsilon'[\ell,k] \neq \Upsilon^*[\ell,k]$ for some $\ell \in [m-1], k \in [\delta-1]$. In this case we will specify two transitions of the heat bath chain, $\Upsilon' \to \Upsilon''$ and $\Upsilon'' \to \Upsilon'''$, so that $d(\Upsilon''', \Upsilon^*) < d(\Upsilon', \Upsilon^*)$.

Assume without loss of generality that $\Upsilon'[\ell,k] > \Upsilon^*[\ell,k]$. Therefore there must be some $k' \in [\delta-1]$ such that $\Upsilon'[\ell,k'] < \Upsilon^*[\ell,k']$.

Now because we only change $Q[i,j]$ values on the path from Υ to Υ^*, $\Upsilon'[\ell,k] > \Upsilon^*[\ell,k]$ implies $\Upsilon'[\ell,k] \geq \Upsilon^*[\ell,k] + (\alpha_{\ell,k}+1)$, and $\Upsilon'[\ell,k'] \leq \Upsilon^*[\ell,k'] + (\alpha_{\ell,k'}+1)$.

By lemma 10 and $\Upsilon'[m,\delta] = \Upsilon^*[m,\delta]$ we know $\Upsilon'[m,\delta] \geq (\alpha_{\ell,k}+1)$. Therefore we can perform the transition $(\Upsilon' \to \Upsilon'')$, where Υ'' is identical to Υ' except for the following entries:

$$\begin{array}{ll} \Upsilon''[\ell,k] = \Upsilon'[\ell,k] - (\alpha_{\ell,k}+1) & \Upsilon''[\ell,\delta] = \Upsilon'[\ell,\delta] + (\alpha_{\ell,k}+1) \\ \Upsilon''[m,k] = \Upsilon'[m,k] + (\alpha_{\ell,k}+1) & \Upsilon''[m,\delta] = \Upsilon'[m,\delta] - (\alpha_{\ell,k}+1) \end{array}$$

By lemma 10 we know $\Upsilon'[\ell,\delta] \geq (\alpha_{\ell,k'}+1)$ and $\Upsilon'[m,k'] \geq (\alpha_{\ell,k'}+1)$. Then we can perform the transition $(\Upsilon'' \to \Upsilon''')$ by changing the following entries:

$$\begin{array}{ll} \Upsilon'''[\ell,k'] = \Upsilon''[\ell,k'] + (\alpha_{\ell,k'}+1) & \Upsilon'''[\ell,\delta] = \Upsilon''[\ell,\delta] - (\alpha_{\ell,k'}+1) \\ \Upsilon'''[m,k'] = \Upsilon''[m,k'] - (\alpha_{\ell,k'}+1) & \Upsilon'''[m,\delta] = \Upsilon''[m,\delta] + (\alpha_{\ell,k'}+1) \end{array}$$

Now we show $d(\Upsilon''', \Upsilon^*) < d(\Upsilon', \Upsilon^*)$.

$$\begin{aligned}
d_1(\Upsilon''', \Upsilon^*) &= d_1(\Upsilon', \Upsilon^*) - (\alpha_{\ell,k} + \alpha_{\ell,k'} + 2) \\
d_2(\Upsilon''', \Upsilon^*) &= d_2(\Upsilon', \Upsilon^*) + \tfrac{N_W+1}{3N_W}|\alpha_{\ell,k} - \alpha_{\ell,k'}| \\
d_3(\Upsilon''', \Upsilon^*) &= d_3(\Upsilon', \Upsilon^*) + \tfrac{N_W+1}{3N_W}(\alpha_{\ell,k} + \alpha_{\ell,k'} + 2) \\
d_4(\Upsilon''', \Upsilon^*) &= d_4(\Upsilon', \Upsilon^*) + \tfrac{N_W+1}{3N_W}|\alpha_{\ell,k} - \alpha_{\ell,k'}|.
\end{aligned}$$

Therefore

$$\begin{aligned}
d(\Upsilon''', \Upsilon^*) &= d(\Upsilon', \Upsilon^*) - (2/3 - 1/3N_W)(\alpha_{\ell,k} + \alpha_{\ell,k'} + 2) + (2/3 + 2/3N_W)|\alpha_{\ell,k} - \alpha_{\ell,k'}| \\
&\leq d(\Upsilon', \Upsilon^*) - (2/3 - 1/3N_W)(\alpha_{\ell,k} + \alpha_{\ell,k'} + 2) + (2/3 + 2/3N_W)(\alpha_{\ell,k} + \alpha_{\ell,k'}) \\
&\leq d(\Upsilon', \Upsilon^*) + (\alpha_{\ell,k} + \alpha_{\ell,k'})/N_W - (2 - 1/N_W)(2/3) \\
&\leq d(\Upsilon', \Upsilon^*) + 2/m^2\delta^2 - (2/3) \\
&< d(\Upsilon', \Upsilon^*).
\end{aligned}$$

4.2.2 Analysis of Flow for Phase 1

The definition of $\alpha_{i,j}$ defines an equivalence class on the set $\Sigma_{\rho,\zeta}$, where $\Phi \equiv \Phi'$ if and only if $\Phi[i,j] = \Phi'[i,j] \bmod (\alpha_{i,j} + 1)$ for every $i \in [m-1], j \in [\delta - 1]$ (i.e. all the remainders are the same).

Note that by definition of the $\alpha_{i,j}$ values, and since $\Phi[i,j] \leq \{\rho_i, \zeta_j\}$ for all $i \in [m-1], j \in [\delta-1]$ for $\Phi \in \Sigma_{\rho,\zeta}$, any equivalence class contains at most $(m^2\delta^2)^{m\delta}$ contingency tables (there are at most $m^2\delta^2$ choices for $Q[i,j]$ for any $i \in [m-1], j \in [\delta-1]$). Therefore each equivalence class contains a constant number of contingency tables.

The routing scheme (a)-(d) defines a path $\Upsilon = \Phi_0, \Phi_1, \ldots, \Phi_t = \Upsilon^*$ from Υ to Υ^*, for every $\Upsilon \in \Sigma_{\rho,\zeta}$. We know Φ_h lies in the same equivalence class as Υ and Υ^* for every h. By our analysis of cases (a)-(d), we know that for every $h \geq 0$, either $d(\Phi_{h+1}, \Upsilon^*) < d(\Phi_h, \Upsilon^*)$ or $d(\Phi_{h+2}, \Upsilon^*) < d(\Phi_h, \Upsilon^*)$. Therefore we can define a subsequence of the path $\{\Phi_h\}$ such that (i) the subsequence contains at least every second element of $\{\Phi_h\}$, and (ii) no contingency table ever appears twice in the subsequence. Therefore the length of the path from Υ to Υ^* is at most $2(m^2\delta^2)^{m\delta}$.

To analyse the amount of flow from Stage 1 that may pass through $\Phi \in \Sigma_{\rho,\zeta}$, we rely on the fact that for any $\Upsilon \in \Sigma_{\rho,\zeta}$ the path from Υ to Υ^* lies in the equivalence class of Υ. Therefore, for any fixed Ψ, there are at most $(m^2\delta^2)^{m\delta}$ different contingency tables Υ that may pass through Φ on the way to Υ^*. Also, by our bound on the length of the path, we know that for any fixed Ψ and Υ, Φ may occur at most $(m^2\delta^2)^{m\delta}$ times on the path from Υ to Υ^*.

Therefore for any fixed Ψ, the flow through any Φ during phase 1 is at most $(m^2\delta^2)^{2m\delta}$. Therefore the total flow through Φ during phase 1 is at most $|\Sigma_{\rho,\zeta}|(m^2\delta^2)^{2m\delta}$.

4.2.3 Phase 2

In this phase we describe how to route Υ^* to Ψ^*. We know that

$$\begin{aligned}
\Upsilon^*[i,j] &= Q^*[i,j](\alpha_{i,j} + 1) + R_\Upsilon[i,j] \\
\Psi^*[i,j] &= Q^*[i,j](\alpha_{i,j} + 1) + R_\Psi[i,j]
\end{aligned}$$

for all $i \in [m-1], j \in [m-1]$, where $Q^*[i,j]$ was defined in (15).

In this phase we route Υ^* to Ψ^* in $m\delta$ steps, by "fixing" one remainder at a time. The key to this approach is part (i) of lemma 10, which shows that for *any* remainders $R[i,j]$ satisfying $R[i,j] \leq \alpha_{i,j}$ for $i \in [m-1], j \in [\delta-1]$, if Φ is defined by

$$\Phi[i,j] = Q^*[i,j](\alpha_{i,j} + 1) + R[i,j]$$

for $i \in [m-1], j \in [\delta-1]$, then $\Phi \in \Sigma_{\rho,\zeta}$.

Suppose we order the "boxes" of the $m \times \delta$ contingency tables in lexicographic order: $(1,1), (1,2), \ldots, (1,\delta-1), (2,1), \ldots, (m-1,1), \ldots, (m-1,\delta-1)$. Let $h = (h_1, h_2)$ denote any point in this lexicographic order. Let h^+ denote the successor to h in this ordering.

Then we can define a series of tables

$$\Upsilon^* = \Phi_{(1,1)}, \Phi_{(1,2)}, \ldots, \Phi_h, \ldots, \Phi_{(m-1,\delta-1)} = \Psi^*$$

by

$$\Phi_h[i,j] = \begin{cases} \Psi^*[i,j] & \text{if } (i,j) \text{ is less than or equal to } h \\ \Upsilon^*[i,j] & \text{otherwise} \end{cases}$$

By part (i) of lemma 10 we know that $\Phi_h \in \Sigma_{\rho,\zeta}$ for all h, and therefore $\Phi_h \to \Phi_{h^+}$ is a transition of the 2×2 heat-bath chain.

Note that if we define a dual table $\bar{\Phi}_h$ by

$$\bar{\Phi}_h[i,j] = \begin{cases} \Upsilon^*[i,j] & \text{if } (i,j) \text{ is less than or equal to } h \\ \Psi^*[i,j] & \text{otherwise} \end{cases}$$

then $\bar{\Phi}_h \in \Sigma_{\rho,\zeta}$ also.

Therefore we have a path of length $(m-1)(\delta-1)$ connecting Υ^* to Ψ^* in $\Sigma_{\rho,\zeta}$.

4.2.4 Analysis of Flow for Phase 2

Now we bound the amount of flow from Phase 2 that can pass through any $\Phi \in \Sigma_{\rho,\zeta}$. We do this using an encoding. Suppose that we are given

(1) A pair of indices $h = (h_1, h_2)$ specifying that Φ occurs as Φ_h on the path from Υ^* to Ψ^* during phase 2;

(2) The dual contingency table $\bar{\Phi}_h$.

(3) The integers $Q_\Upsilon[i,j]$ for all $i \in [m], j \in [\delta]$.

(4) The integers $Q_\Psi[i,j]$ for all $i \in [m], j \in [\delta]$.

Then we can construct the original pair of tables Υ and Ψ exactly. The information in (1) and (2) first allows us to reconstruct Υ^* and Ψ^*. Then using Υ^*, we may reconstruct $R_\Upsilon[i,j]$ for all i,j since we know (or may compute) $Q^*[i,j]$. Knowing $R_\Upsilon[i,j]$, together with the information in (3), we can find Υ exactly, and in a like manner we can reconstruct Ψ. There are at most $(m^2\delta^2)^{m\delta}$ possible values for the $Q_\Upsilon[i,j]$, and the same number for the $Q_\Psi[i,j]$.

Therefore, for any $\Phi \in \Sigma_{\rho,\zeta}$, the total amount of flow that may pass through Φ during phase 2 is at most $(m\delta)(m^2\delta^2)^{2m\delta}|\Sigma_{\rho,\zeta}|$.

4.2.5 Finishing up

Combining Phase 1 and 2, the length of any (Υ, Ψ)-path is at most $4(m\delta)^{2m\delta} + m\delta$. The total amount of flow that can pass through any $\Phi \in \Sigma_{\rho,\zeta}$ is at most $|\Sigma_{\rho,\zeta}|(2(m\delta)^{4m\delta} + m\delta(m^2\delta^2)^{2m\delta})$.

This establishes the condition on the flow f^* that was cited in section 4.1. We then alter this flow as outlined in that section to get the new flow f, letting us establish theorem 9, proving rapid mixing of the 2×2 heat-bath chain on the set of $m \times n$ contingency tables $\Sigma_{r,c}$.

References

[1] F.K.R Chung, R.L. Graham and S.-T. Yau, On sampling with Markov chains. *Random Structures & Algorithms*, **9**(1-2), 1996, pp. 55–77.

[2] M. Cryan and M. Dyer, A polynomial time algorithm to approximately count contingency tables when the number of rows is constant, to appear in *Proceedings of the 34th ACM Symposium on Theory of Computing*, Montréal, 2002.

[3] J.A. De Loera and B. Sturmfels, Algebraic unimodular counting, Preprint, Department of Mathematics, University of California, Davis, 2001.

[4] P. Diaconis and B. Efron, Testing for independence in a two-way table: new interpretations of the chi-square statistic (with discussion). *Annals of Statistics*, **13**, 1995, pp. 845–913.

[5] P. Diaconis and A. Gangolli, Rectangular arrays with fixed margins, in: D. Aldous, P.P. Varaiya, J. Spencer and J.M. Steele (Eds.), *Discrete Probability and Algorithms*, IMA Volumes on Mathematics and its Applications, **72**, Springer, New York, 1995, pp. 15–41.

[6] P. Diaconis and L. Saloff-Coste, Comparison theorems for reversible Markov chains. *The Annals of Applied Probability*, **3**(3), 1993, pp. 696–730.

[7] P. Diaconis and L. Saloff-Coste, Random walk on contingency tables with fixed row and column sums. Technical Report, Department of Mathematics, Harvard University, 1995.

[8] P. Diaconis and D. Stroock, Geometric bounds for eigenvalues of Markov chains. *The Annals of Applied Probability*, **1**, 1991, pp. 36–61.

[9] M. Dyer and C. Greenhill, Polynomial-time counting and sampling of two-rowed contingency tables. *Theoretical Computer Science*, **246**, 2000, pp. 265–278.

[10] M. Dyer, R. Kannan and J. Mount, Sampling contingency tables. *Random Structures & Algorithms*, **10**(4), 1997, pp. 487–506.

[11] G.R. Grimmett and D.R. Stirzaker, *Probability and Random Processes*, Oxford University Press, Oxford, 1992.

[12] V.S. Grinberg and S.V. Sevast'yanov, Value of the Steinitz constant. *Funktsional. Anal. i Prilozhen.*, **14**(2), 1980, pp. 56–57.

[13] D. Hernek, Random generation of $2 \times n$ contingency tables. *Random Structures & Algorithms*, **13**(1), 1998, pp. 71–79.

[14] M. Jerrum and A. Sinclair, Markov chain Monte Carlo method: an approach to approximate counting and integration. D.S. Hochbaum (Ed.), *Approximation Algorithms for NP-Hard Problems*, PWS, Boston, 1997, pp. 482–520.

[15] M.R. Jerrum, L.G. Valiant, and V.V. Vazirani, Random generation of combinatorial structures from a uniform distribution, *Theoretical Computer Science,* **43**, 1986, pp. 169–188.

[16] B. Morris, Improved bounds for sampling contingency tables. *3rd International Workshop on Randomization and Approximation Techniques in Computer Science*, volume 1671 of *Lecture Notes in Computer Science*, 1999, pp. 121–129.

[17] B. Morris, *Random Walks in Convex Sets*. PhD thesis, Department of Statistics, University of California, Berkeley, 2000.

[18] B. Morris and A.J. Sinclair, Random walks on truncated cubes and sampling 0-1 knapsack solutions. In *40th IEEE Symposium on Foundations of Computer Science*, 1999, pp. 230–240.

[19] J. Mount, *Application of convex sampling to optimization and contingency table generation.* PhD thesis, Technical report CMU-CS-95-152, Computer Science Department, Carnegie Mellon University, 1995.

[20] I. Pak, On sampling integer points in polyhedra, Preprint, Department of Mathematics, Yale University, 2000.

[21] D. Randall and P. Tetali, Analyzing Glauber dynamics by comparison of Markov chains. *Journal of Mathematical Physics* **41**(3), 2000, pp. 1598–1615.

[22] A.J. Sinclair, Improved bounds for mixing rates of Markov chains and multicommodity flow. *Combinatorics, Probability and Computing*, **1**, 1992, pp. 351–370.

[23] E. Steinitz, bedingt konvergente Reihen und konvexe Systeme, *J. reine angew. Math.*, **143**, 1913, pp. 128–175.

[24] E. Vigoda, Improved bounds for sampling colorings. *Journal of Mathematical Physics* **41**(3), 2000, pp. 1555–1569.